

Rabbit anti-Human MyD88 Monoclonal Antibody | anti-MYD88 antibody

[KO Validated] MyD88 Rabbit PolymAb

CATALOG # AAA282976	HOST Rabbit	REACTIVITY Human	APPLICATIONS ELISA, Western Blot
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

MyD88; Monoclonal Antibody; [KO Validated] MyD88 Rabbit PolymAb; WM1; IMD68; MYD88D; anti-MYD88 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.05% proclin300, 0.05% BSA, 50% glycerol, pH7.3.
Sequence	LAEEMDFEYL EIRQLETQAD PTGRLLDAWQ GRPGASVGRL LELLTKLGRD DVLLELGPSI EEDCQKYILK QQQEEAEKPL QVAAVDSSVP RTAELAGITT L
Applicable Applications for anti-MYD88 antibody	ELISA, WB (Western Blot)
Cross Reactivity	Human, Mouse, Rat
Immunogen	A synthetic peptide corresponding to a sequence within amino acids 50-150 of human MyD88 (NP_002459.3).
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES



WB (Western Blot)

Western blot analysis of various lysates, using MyD88 Rabbit PolymAb® (AAA282976) at 1:8000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Exposure time: 60s.



WB (Western Blot)

Western blot analysis of various lysates, using MyD88 Rabbit PolymAb® (AAA282976) at 1:8000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 30s.

ADDITIONAL INFORMATION

Related Product Information for anti-MYD88 antibody	This gene encodes a cytosolic adapter protein that plays a central role in the innate and adaptive immune response. This protein functions as an essential signal transducer in the interleukin-1 and Toll-like receptor signaling pathways. These pathways regulate that activation of numerous proinflammatory genes. The encoded protein consists of an N-terminal death domain and a C-terminal Toll-interleukin1 receptor domain. Patients with defects in this gene have an increased susceptibility to pyogenic bacterial infections. Alternate splicing results in multiple transcript variants.
Product Categories/Family for anti-MYD88 antibody	Signal Transduction; Immunology Inflammation; NF-kB Signaling Pathway; Toll-like Receptor Signaling Pathway; Cell Intrinsic Innate Immunity Signaling Pathway; TLR Signaling; Cardiovascular; Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	289546500
NCBI GeneID	4615
NCBI Accession #	NP_001166037.1
NCBI GenBank Nucleotide #	NM_001172566.1
UniProt Accession #	Q99836; P78397; Q53XS7; B4DKH8; B4DQ60; B4DQ72
Molecular Weight	Calculated MW: 33kDa Observed MW: 33kDa
UniProt Protein Name	Myeloid differentiation primary response protein MyD88
UniProt Gene Name	MYD88
UniProt Entry Name	MYD88_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/monoclonal/myd88-antibody-western-blot-elisa/282976>