

Rabbit anti-Human COL1A1 Polyclonal Antibody | anti-COL1A1 antibody

COL1A1 Rabbit pAb

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA283300	Rabbit	Human	ELISA, Immunocytochemistry, Immunofluorescence, Western Blot


PRODUCT SYNONYMS

COL1A1; Polyclonal Antibody; COL1A1 Rabbit pAb; OI1; OI2; OI3; OI4; EDSC; CAFYD; EDSARTH1; Collagen I/COL1A1; anti-COL1A1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.05% proclin300, 50% glycerol, pH7.3.
Sequence	STEASQNITY HCKNSVAYMD QQTGNLKKAL LLQGSNEIEI RAEGNSRFTY SVTVDGCTSH TGAWGKTVIE YKTTKTSRLP IIDVAPLDVG APDQEFQFDV GPCVFL
Applicable Applications for anti-COL1A1 antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), WB (Western Blot)
Cross Reactivity	Human, Mouse
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 1359-1464 of human COL1A1 (NP_000079.2).
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES



IF (Immunofluorescence)

Immunofluorescence analysis of NIH/3T3 cells using COL1A1 Rabbit pAb(AAA283300) at a dilution of 1:250 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L)(AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



WB (Western Blot)

Western blot analysis of various lysates using COL1A1 Rabbit pAb (AAA283300) at 1:3000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Negative control (NC): HT-29. Exposure time: 30s.

ADDITIONAL INFORMATION

Related Product Information for anti-COL1A1 antibody	This gene encodes the pro-alpha1 chains of type I collagen whose triple helix comprises two alpha1 chains and one alpha2 chain. Type I is a fibril-forming collagen found in most connective tissues and is abundant in bone, cornea, dermis and tendon. Mutations in this gene are associated with osteogenesis imperfecta types I-IV, Ehlers-Danlos syndrome type VIIA, Ehlers-Danlos syndrome Classical type, Caffey Disease and idiopathic osteoporosis. Reciprocal translocations between chromosomes 17 and 22, where this gene and the gene for platelet-derived growth factor beta are located, are associated with a particular type of skin tumor called dermatofibrosarcoma protuberans, resulting from unregulated expression of the growth factor. Two transcripts, resulting from the use of alternate polyadenylation signals, have been identified for this gene.
Product Categories/Family for anti-COL1A1 antibody	Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	296439504
NCBI GeneID	1277
UniProt Accession #	P02452; O76045; P78441; Q13896; Q13902; Q13903; Q14037; Q14992; Q15176; Q15201; Q16050; Q59F64
Molecular Weight	Calculated MW: 139kDa Observed MW: 220kDa
UniProt Protein Name	Collagen alpha-1(I) chain
UniProt Gene Name	COL1A1
UniProt Entry Name	CO1A1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/col1a1-antibody-immunocytochemistry-immunofluorescence-western-blot-elisa/283300>