

Rabbit anti-Human ABCC5 Polyclonal Antibody | anti-ABCC5 antibody

ABCC5 Antibody (Center)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA284077	Rabbit	Human	Flow Cytometry, Functional Assay, Immunohistochemistry, ELISA, Western Blot


PRODUCT SYNONYMS

ABCC5; Polyclonal Antibody; ABCC5 Antibody (Center); Multidrug resistance-associated protein 5; ATP-binding cassette sub-family C member 5; Multi-specific organic anion transporter C; MOAT-C; SMRP; pABC11; MRP5; anti-ABCC5 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	This ABCC5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 625-652 amino acids from the Central region of human ABCC5.
Purity/Purification	This antibody is purified through a protein A column, followed by peptide affinity purification.
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Applicable Applications for anti-ABCC5 antibody	FCM/FACS (Flow Cytometry), IHC (Immunohistochemistry), ELISA, WB (Western Blot)
Antigen Region	625-652
Function	ATP-dependent transporter of the ATP-binding cassette (ABC) family that actively extrudes physiological compounds, and xenobiotics from cells. Mediates ATP-dependent transport of endogenous metabolites such as cAMP and cGMP, folic acid and N-lactoyl-amino acids (in vitro) (PubMed:10893247, PubMed:12637526, PubMed:12695538, PubMed:15899835, PubMed:17229149, PubMed:25964343). Acts also as a general glutamate conjugate and analog transporter that can limit the brain levels of endogenous metabolites, drugs, and toxins (PubMed:26515061). Confers resistance to the antiviral agent PMEA (PubMed:12695538). Able to transport several anticancer drugs including methotrexate, and nucleotide analogs in vitro, however it does with low affinity, thus the exact role of ABCC5 in mediating resistance still needs to be elucidated (PubMed:10840050, PubMed:12435799, PubMed:12695538, PubMed:15899835). Acts as a heme transporter required for the translocation of cytosolic heme to the secretory pathway (PubMed:24836561). May play a role in energy metabolism by regulating the glucagon-like peptide 1 (GLP-1) secretion from enteroendocrine cells (By similarity).
Cellular Location	Basolateral cell membrane; Multi-pass membrane protein. Golgi apparatus lumen Endosome membrane. Cytoplasmic granule {ECO:0000250 UniProtKB:Q9R1X5}. Apical cell membrane; Multi-pass membrane protein. Note=In most cells, routes to the basolateral plasma membrane, but in the brain capillary endothelial cells that form the blood-brain barrier, resides in the apical membrane
Tissue Location	[Isoform 3]: Predominant isoform in retinal pigment epithelium, bladder, and stomach.
Preparation and Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.


IMAGES



FCM/FACS (Flow Cytometry)
Flow cytometric analysis of 293 cells using ABCC5 Antibody (Center)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



IHC (Immunohistochemistry)
Formalin-fixed and paraffin-embedded human brain tissue with ABCC5 Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



WB (Western Blot)
Western blot analysis of ABCC5 Antibody (Center) (AAA284077) in 293 cell line lysates (35ug/lane). ABCC5 (arrow) was detected using the purified Pab.(2ug/ml)

ADDITIONAL INFORMATION

Related Product Information for anti-ABCC5 antibody	ABCC5 is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the MRP subfamily which is involved in multi-drug resistance. This protein functions in the cellular export of its substrate, cyclic nucleotides. This export contributes to the degradation of phosphodiesterases and possibly an elimination pathway for cyclic nucleotides. Studies show that this protein provides resistance to thiopurine anticancer drugs,6-mercaptopurine and thioguanine, and the anti-HIV drug 9-(2-phosphonylmethoxyethyl)adenine. This protein may be involved in resistance to thiopurines in acute lymphoblastic leukemia and antiretroviral nucleoside analogs in HIV-infected patients.
Product Categories/Family for anti-ABCC5 antibody	Signal Transduction
References	Olsen,J.V., et.al., Cell 127 (3), 635-648 (2006)

NCBI AND UNIPROT INFORMATION

NCBI GI #	66529093
NCBI GeneID	10057
NCBI Accession #	NP_001018881.1; NP_005679.2

NCBI GenBank Nucleotide #	NM_001023587.1
UniProt Accession #	O15440 ; O14517 ; Q29ZA9 ; Q29ZB1 ; Q86UX3 ; Q86W30 ; Q9UN85 ; Q9UNP5 ; Q9UQC3 ; B9EIQ2
Molecular Weight	160660
NCBI Official Full Name	multidrug resistance-associated protein 5 isoform 2
NCBI Official Synonym Full Names	ATP-binding cassette, sub-family C (CFTR/MRP), member 5
NCBI Official Symbol	ABCC5
NCBI Official Synonym Symbols	MRP5; SMRP; ABC33; MOATC; MOAT-C; pABC11; EST277145
NCBI Protein Information	multidrug resistance-associated protein 5
UniProt Protein Name	Multidrug resistance-associated protein 5
UniProt Gene Name	ABCC5
UniProt Synonym Gene Names	MRP5; MOAT-C
UniProt Entry Name	MRP5_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/abcc5-antibody-flow-cytometry-immunohistochemistry-western-blot-elisa/284077>