

Rabbit CFL1 Polyclonal Antibody | anti-CFL1 antibody

CFL1 Antibody (Center)


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA285656	Rabbit	Human, mouse (Predicted Reactivity: Bovine, Monkey, Pig, Rat, Chicken, Xenopus)	Flow Cytometry, Functional Assay, ELISA, Western Blot

PRODUCT SYNONYMS


CFL1; Polyclonal Antibody; CFL1 Antibody (Center); Cofilin-1; 18 kDa phosphoprotein; p18; Cofilin; non-muscle isoform; CFL; anti-CFL1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, mouse (Predicted Reactivity: Bovine, Monkey, Pig, Rat, Chicken, Xenopus)
Clonality	Polyclonal
Isotype	Rabbit Ig
Specificity	This CFL1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 70-98 amino acids from the Central region of human CFL1.
Purity/Purification	Peptide Affinity Purified Rabbit Polyclonal Antibody (Pab)
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Sequence Positions	70-98
Sequence Length	166
Applicable Applications for anti-CFL1 antibody	FCM/FACS (Flow Cytometry), ELISA, WB (Western Blot)
Antigen Type	Synthetic Peptide
Antigen Source	HUMAN
Function	Binds to F-actin and exhibits pH-sensitive F-actin depolymerizing activity (PubMed:11812157). In conjunction with the subcortical maternal complex (SCMC), plays an essential role for zygotes to progress beyond the first embryonic cell divisions via regulation of actin dynamics (PubMed:15580268). Required for the centralization of the mitotic spindle and symmetric division of zygotes (By similarity). Plays a role in the regulation of cell morphology and cytoskeletal organization in epithelial cells (PubMed:21834987). Required for the up-regulation of atypical chemokine receptor ACKR2 from endosomal compartment to cell membrane, increasing its efficiency in chemokine uptake and degradation (PubMed:23633677). Required for neural tube morphogenesis and neural crest cell migration (By similarity).
Cellular Location	Nucleus matrix. Cytoplasm, cytoskeleton. Cell projection, ruffle membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, lamellipodium membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, lamellipodium {ECO:0000250 UniProtKB:P18760}. Note=Colocalizes with the actin cytoskeleton in membrane ruffles and lamellipodia. Detected at the cleavage furrow and contractile ring during cytokinesis. Almost completely in nucleus in cells exposed to heat shock or 10% dimethyl sulfoxide
Tissue Location	Widely distributed in various tissues.
Preparation and Storage	Maintain refrigerated at 2-8°C. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.




WB (Western Blot)
Anti-CFL1 Antibody (Center) at 1:1000 dilution + MCF-7 whole cell lysate Lysates/proteins at 20 ug per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 19 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



FCM/FACS (Flow Cytometry)
CFL1 Antibody (Center) (AAA285656) flow cytometric analysis of HL-60 cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



WB (Western Blot)
Western blot analysis of CFL1 (arrow) using rabbit polyclonal CFL1 Antibody (Center) (AAA285656). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the CFL1 gene.



WB (Western Blot)
Western blot analysis of CFL1 Antibody (Center) (AAA285656) in HL-60 cell line lysates (35ug/lane). CFL1 (arrow) was detected using the purified Pab;

ADDITIONAL INFORMATION

Related Product Information for anti-CFL1 antibody	Cofilin is a widely distributed intracellular actin-modulating protein that binds and depolymerizes filamentous F-actin and inhibits the polymerization of monomeric G-actin in a pH-dependent manner.
Product Categories/Family for anti-CFL1 antibody	Cancer; Cardiovascular; Neuroscience; Signal transduction
References	Fazal,F., et.al., J. Biol. Chem. 284 (31), 21047-21056 (2009)

NCBI AND UNIPROT INFORMATION

NCBI GI #	5031635
NCBI GeneID	1072
NCBI Accession #	NP_005498.1
NCBI GenBank Nucleotide #	NM_005507.2
UniProt Accession #	P23528; Q53Y87; Q9UCA2; B3KUQ1
Molecular Weight	18502
NCBI Official Full Name	cofilin-1

NCBI Official Synonym Full Names	cofilin 1 (non-muscle)
NCBI Official Symbol	CFL1
NCBI Official Synonym Symbols	CFL; cofilin; HEL-S-15
NCBI Protein Information	cofilin-1
UniProt Protein Name	Cofilin-1
UniProt Gene Name	CFL1
UniProt Synonym Gene Names	CFL; p18
UniProt Entry Name	COF1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/cfl1-antibody-flow-cytometry-western-blot-elisa/285656>