

Rabbit anti-Human, mouse MARK1 Polyclonal Antibody | anti-MARK1 antibody

MARK1 Antibody (N-term)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA286132	Rabbit	Human, mouse	Western Blot, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

MARK1; Polyclonal Antibody; MARK1 Antibody (N-term); Serine/threonine-protein kinase MARK1; MAP/microtubule affinity-regulating kinase 1; PAR1 homolog c; Par-1c; Par1c; MARK1 HGNC:6896; anti-MARK1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, mouse
Clonality	Polyclonal
Isotype	Rabbit Ig
Specificity	This MARK1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 6-40 amino acids from the N-terminal region of human MARK1.
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Concentration	Vial Concentration: 2 (varies by lot)
Sequence Positions	6-40
Sequence Length	758
Applicable Applications for anti-MARK1 antibody	WB (Western Blot), IHC (Immunohistochemistry), ELISA
Antigen Type	Synthetic Peptide
Antigen Source	HUMAN
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 6 months. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



IHC (Immunohiostchemistry)
Formalin-fixed and paraffin-embedded human brain tissue reacted with MARK1 Antibody (N-term) , which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



WB (Western Blot)
The anti-MARK1 N-term Pab is used in Western blot to detect MARK1 in P7 mouse whole brain lysate (60 ug). 1:250 dilution of anti-MARK1 pab was used. Data and protocol kindly provided by DR. Shengli Zhao, M.D., Ph.D. of Duke University Medical Center.

ADDITIONAL INFORMATION

Related Product Information for anti-MARK1 antibody	MARK is a family of kinases that is known for its involvement in establishing cell polarity and in phosphorylating tau protein during Alzheimer neurodegeneration. Expression of MARK causes the phosphorylation of MAPs at their KXGS motifs, thereby detaching MAPs from the microtubules and thus facilitating the transport of particles. This occurs without impairing the intrinsic activity of motors because the velocity during active movement remains unchanged. In primary retinal ganglion cells, transfection with tau leads to the inhibition of axonal transport of mitochondria, APP vesicles, and other cell components which leads to starvation of axons and vulnerability against stress. This transport inhibition can be rescued by phosphorylating tau with MARK
Product Categories/Family for anti-MARK1 antibody	Signal Transduction
References	Drewes, G., et al., Cell 89(2):297-308 (1997).

NCBI AND UNIPROT INFORMATION

NCBI GI #	153791472
NCBI GeneID	4139
NCBI Accession #	NP_061120.3
NCBI GenBank Nucleotide #	NM_018650.4
UniProt Accession #	Q9P0L2 ; Q2HIY1 ; Q5VTF9 ; Q5VTG0 ; Q96SW9 ; Q9P251 ; D3DTB0 ; D3DTB1
Molecular Weight	89003
NCBI Official Full Name	serine/threonine-protein kinase MARK1 isoform 2
NCBI Official Synonym Full Names	MAP/microtubule affinity-regulating kinase 1
NCBI Official Symbol	MARK1
NCBI Official Synonym Symbols	MARK; Par1c; Par-1c
NCBI Protein Information	serine/threonine-protein kinase MARK1
UniProt Protein Name	Serine/threonine-protein kinase MARK1
UniProt Gene Name	MARK1
UniProt Synonym Gene Names	Par-1c; Par1c

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/mark1-antibody-western-blot-immunohistochemistry-elisa/286132>