

Rabbit anti-Human NFATC4 Polyclonal Antibody | anti-NFATC4 antibody

NFATC4 Antibody (C-term)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA286532	Rabbit	Human	Western Blot, ELISA, Immunofluorescence


PRODUCT SYNONYMS

NFATC4; Polyclonal Antibody; NFATC4 Antibody (C-term); Nuclear factor of activated T-cells; cytoplasmic 4; NF-ATc4; NFATc4; T-cell transcription factor NFAT3; NF-AT3; NFAT3; anti-NFATC4 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit Ig
Specificity	This NFATC4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 744-773 amino acids from the C-terminal region of human NFATC4.
Purity/Purification	Peptide Affinity Purified Rabbit Polyclonal Antibody (Pab)
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Concentration	Vial Concentration: 0.5 (varies by lot)
Sequence Positions	744-773
Sequence Length	964
Applicable Applications for anti-NFATC4 antibody	WB (Western Blot), ELISA, IF (Immunofluorescence)
Antigen Type	Synthetic Peptide
Crown Antibody	Yes
Antigen Source	HUMAN
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 6 months. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



IF (Immunofluorescence)
Fluorescent confocal image of U251 cell stained with NFATC4 Antibody (C-term). U251 cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.1%, 10 min), then incubated with NFATC4 primary antibody (1:25, 1 h at 37 degree). For secondary antibody, Alexa Fluor 488 conjugated donkey anti-rabbit antibody (green) was used (1:400, 50 min at 37 degree). Cytoplasmic actin was counterstained with Alexa Fluor 555 (red) conjugated Phalloidin (7units/ml, 1 h at 37 degree). Nuclei were counterstained with DAPI (blue) (10 ug/ml, 10 min). NFATC4 immunoreactivity is localized to Cytoplasm significantly.



WB (Western Blot)
NFATC4 Antibody (C-term) western blot analysis in human placenta tissue lysates (35ug/lane). This demonstrates the NFATC4 antibody detected the NFATC4 protein (arrow).

ADDITIONAL INFORMATION

Related Product Information for anti-NFATC4 antibody	The product of this gene is a member of the nuclear factors of activated T cells DNA-binding transcription complex. This complex consists of at least two components: a preexisting cytosolic component that translocates to the nucleus upon T cell receptor (TCR) stimulation and an inducible nuclear component. Other members of this family of nuclear factors of activated T cells also participate in the formation of this complex. The product of this gene plays a role in the inducible expression of cytokine genes in T cells, especially in the induction of the IL-2 and IL-4. Alternatively spliced transcript variants encoding different isoforms have been noted for this gene. [provided by RefSeq].
Product Categories/Family for anti-NFATC4 antibody	Crown Antibodies; Immunology; Signal Transduction
References	Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010) Al-Daraji, W.I., et al. J. Cutan. Pathol. 37 (9), E21-E36 (2010) : Fougere, M., et al. Oncogene 29(15):2292-2301(2010) Davila, S., et al. Genes Immun. 11(3):232-238(2010) Talmud, P.J., et al. Am. J. Hum. Genet. 85(5):628-642(2009)

NCBI AND UNIPROT INFORMATION

NCBI GI #	209862845
NCBI GeneID	4776
NCBI Accession #	NP_001129494.1; NP_001185894.1; NP_001185895.1; NP_001185896.1; NP_004545.2
NCBI GenBank Nucleotide #	NM_001136022.2
UniProt Accession #	Q14934; B4DDG5; B4DY55; B5B2U7; B5B2U8; B5B2U9; B5B2V0; B5B2V1; B5B2V2; B5B2V3; B5B2V4; B5B2V5
Molecular Weight	95449
NCBI Official Full Name	nuclear factor of activated T-cells, cytoplasmic 4 isoform 1
NCBI Official Synonym Full Names	nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 4
NCBI Official Symbol	NFATC4
NCBI Official Synonym Symbols	NFAT3; NF-AT3; NF-ATC4
NCBI Protein Information	nuclear factor of activated T-cells, cytoplasmic 4

UniProt Protein Name	Nuclear factor of activated T-cells, cytoplasmic 4
UniProt Gene Name	NFATC4
UniProt Synonym Gene Names	NFAT3; NF-ATc4; NFATc4; NF-AT3
UniProt Entry Name	NFAC4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/nfatc4-antibody-western-blot-immunofluorescence-elisa/286532>