

Rabbit CD38 Polyclonal Antibody | anti-CD38 antibody

CD38 Antibody (C-term)


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA28702	Rabbit	Human (Predicted Reactivity: Monkey)	Western Blot, ELISA, Immunohistochemistry, Immunofluorescence, Flow Cytometry, Functional Assay

PRODUCT SYNONYMS

CD38; Polyclonal Antibody; CD38 Antibody (C-term); ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1; 2'-phospho-ADP-ribosyl cyclase; 2'-phospho-ADP-ribosyl cyclase/2'-phospho-cyclic-ADP-ribose transferase; 2'-phospho-cyclic-ADP-ribose transferase; ADP-ribosyl cyclase 1; ADPRC 1; Cyclic ADP-ribose hydrolase 1; cADPr hydrolase 1; T10; anti-CD38 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human (Predicted Reactivity: Monkey)
Clonality	Polyclonal
Isotype	Rabbit Ig
Specificity	This CD38 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 241-270 amino acids from the C-terminal region of human CD38.
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Concentration	Vial Concentration: 2 (varies by lot)
Sequence Positions	241-270
Sequence Length	122
Applicable Applications for anti-CD38 antibody	WB (Western Blot), ELISA, IHC (Immunohistochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Application Notes	WB~~1:1000
Antigen Type	Synthetic Peptide
Crown Antibody	Yes
Antigen Source	HUMAN
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 6 months. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.




IF (Immunofluorescence)

Western blot analysis of lysate from RPMI 8226 cell line, using CD38 Antibody (C-term). AAA28702 was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.




IF (Immunofluorescence)

Confocal immunofluorescent analysis of CD38 Antibody (C-term) with Hela cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).




FCM (Flow Cytometry)

Flow cytometric analysis of HepG2 cells using CD38 Antibody (C-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.




IF (Immunofluorescence)

Immunofluorescence analysis of CD38 Antibody (C-term) with paraffin-embedded human hepatocarcinoma tissue. 0.05 mg/ml primary antibody was followed by FITC-conjugated goat anti-rabbit IgG (whole molecule). FITC emits green fluorescence. Red counterstaining is PI.



IHC (Immunohistochemistry)

Formalin-fixed and paraffin-embedded human hepatocarcinoma with CD38 Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



WB (Western Blot)

Western blot analysis of lysate from RPMI8226 cell line, using CD38 Antibody (C-term). AAA28702 was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20ug.

ADDITIONAL INFORMATION

Related Product Information for anti-CD38 antibody

CD38 is a novel multifunctional ectoenzyme widely expressed in cells and tissues especially in leukocytes. CD38 also functions in cell adhesion,signal transduction and calcium signaling.

Product Categories/Family for anti-CD38 antibody	Crown Antibodies ; Cancer ; Immunology ; Neuroscience ; Signal Transduction
References	Horenstein,A.L., Mol. Med. 15 (3-4), 76-84 (2009) Liu,Q., Biochemistry 47 (52), 13966-13973 (2008) Liu,Q., Chem. Biol. 15 (10), 1068-1078 (2008)

NCBI AND UNIPROT INFORMATION

NCBI GI #	38454326
NCBI GeneID	952
NCBI Accession #	NP_001766.2
NCBI GenBank Nucleotide #	NM_001775.2
UniProt Accession #	P28907 ; O00121 ; O00122 ; Q96HY4
Molecular Weight	34328
NCBI Official Full Name	ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1
NCBI Official Synonym Full Names	CD38 molecule
NCBI Official Symbol	CD38
NCBI Official Synonym Symbols	T10; ADPRC 1
NCBI Protein Information	ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1
UniProt Protein Name	ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1
UniProt Gene Name	CD38
UniProt Synonym Gene Names	ADPRC 1; cADPr hydrolase 1
UniProt Entry Name	CD38_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/cd38-antibody-western-blot-immunohistochemistry-immunofluorescence-flow-cytometry-elisa/28702>