

Rabbit anti-Human, mouse EphA4 Polyclonal Antibody | anti-EPHA4 antibody

EphA4 Antibody (C-term)


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA28736	Rabbit	Human, mouse	Western Blot, ELISA, Immunohistochemistry

PRODUCT SYNONYMS


EphA4; Polyclonal Antibody; EphA4 Antibody (C-term); Ephrin type-A receptor 4; EPH-like kinase 8; EK8; hEK8; Tyrosine-protein kinase TYRO1; Tyrosine-protein kinase receptor SEK; EPHA4; HEK8; SEK; TYRO1; anti-EPHA4 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, mouse
Clonality	Polyclonal
Isotype	Rabbit Ig
Specificity	This EphA4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 875-904 amino acids from the C-terminal region of human EphA4.
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Concentration	Vial Concentration: 2 (varies by lot)
Sequence Positions	875-904
Sequence Length	935
Applicable Applications for anti-EPHA4 antibody	WB (Western Blot), ELISA, IHC (Immunohistochemistry)
Application Notes	WB~~1:1000
Antigen Type	Synthetic Peptide
Antigen Source	HUMAN
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 6 months. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.




IHC (Immunohistochemistry)
Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.




WB (Western Blot)
Western blot analysis of EPHA4 (arrow) using EphA4 Antibody (C-term). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the EPHA4 gene (Lane 2) (Origene Technologies).




WB (Western Blot)
Western blot analysis of anti-EphA4 Pab in NCI-H460 cell lysate. EphA4 (arrow) was detected using purified Pab. Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence.



WB (Western Blot)
Western blot analysis of lysates from HeLa, NCI-H460, mouse NIH/3T3 cell line and human ovary tissue lysate (from left to right), using EPHA4 Antibody (R890). AAA28736 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 35ug per lane.



WB (Western Blot)
Western blot analysis of lysates from HeLa, NCI-H460, HUVEC cell line (from left to right), using EPHA4 Antibody (R890). AAA28736 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.



WB (Western Blot)
Western blot analysis of lysates from HeLa, HUVEC cell line (from left to right), using EPHA4 Antibody (R890). AAA28736 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

ADDITIONAL INFORMATION

Related Product Information for anti-EPHA4 antibody	Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the γ phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains. The tyrosine kinase (TK) group is mainly involved in the regulation of cell-cell interactions such as differentiation, adhesion, motility and death. There are currently about 90 TK genes sequenced, 58 are of receptor protein TK (e.g. EGFR, EPH, FGFR, PDGFR, TRK, and VEGFR families), and 32 of cytosolic TK (e.g. ABL, FAK, JAK, and SRC families).
Product Categories/Family for anti-EPHA4 antibody	Cardiovascular; Neuroscience; Signal Transduction
References	Prevost, N., et al., Proc. Natl. Acad. Sci. U.S.A. 99(14):9219-9224 (2002). Xu, Q., et al., Philos. Trans. R. Soc. Lond., B, Biol. Sci. 355(1399):993-1002 (2000). Holder, N., et al., Development 126(10):2033-2044 (1999). Zhou, R., Pharmacol. Ther. 77(3):151-181 (1998). Fox, G.M., et al., Oncogene 10(5):897-905 (1995).

NCBI AND UNIPROT INFORMATION

NCBI GI #	4758280
NCBI GeneID	2043
NCBI Accession #	NP_004429.1
NCBI GenBank Nucleotide #	NM_004438.4
UniProt Accession #	P54764; Q2M380; A8K2P1; B2R601; B7Z6Q8
Molecular Weight	109860
NCBI Official Full Name	ephrin type-A receptor 4 isoform a
NCBI Official Synonym Full Names	EPH receptor A4
NCBI Official Symbol	EPHA4
NCBI Official Synonym Symbols	SEK; HEK8; TYRO1
NCBI Protein Information	ephrin type-A receptor 4
UniProt Protein Name	Ephrin type-A receptor 4
UniProt Gene Name	EPHA4
UniProt Synonym Gene Names	HEK8; SEK; TYRO1; EK8; hEK8
UniProt Entry Name	EPHA4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/epha4-antibody-western-blot-immunohistochemistry-elisa/28736>