

Rabbit anti-Human Cleaved-Caspase-8 Polyclonal Antibody | anti-CASP8 antibody

Cleaved-Caspase-8 (D384) Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA28833	Rabbit	Human	Western Blot, Immunohistochemistry, Immunofluorescence


PRODUCT SYNONYMS

Cleaved-Caspase-8; Polyclonal Antibody; Cleaved-Caspase-8 (D384) Polyclonal Antibody; CASP8; MCH5; Caspase-8; CASP-8; Apoptotic cysteine protease; Apoptotic protease Mch-5; CAP4; FADD-homologous ICE/ced-3-like protease; FADD-like ICE; FLICE; ICE-like apoptotic protease 5; MORT1-associated ced-3 homolog; MACH; anti-CASP8 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Form/Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Applicable Applications for anti-CASP8 antibody	WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence)
Application Notes	WB: 1:500-2000 IF 1:50-300 IHC: 1:50-300
Tissue Location	Isoform 1, isoform 5 and isoform 7 are expressed in a wide variety of tissues. Highest expression in peripheral blood leukocytes, spleen, thymus and liver. Barely detectable in brain, testis and skeletal muscle
Notes	ECO:0000303 PubMed:9931493, ECO:0000312 HGNC: HGNC:1509
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IF (Immunofluorescence)

Dilution: WB 1:500-2000, IF 1:50-300, IHC 1:50-300




IHC (Immunohistochemistry)

Dilution: WB 1:500-2000, IF 1:50-300, IHC 1:50-300




IHC (Immunohistochemistry)

Dilution: WB 1:500-2000, IF 1:50-300, IHC 1:50-300




IHC (Immunohistochemistry-Paraffin)

Dilution: WB 1:500-2000, IF 1:50-300, IHC 1:50-300



IHC (Immunohistochemistry-Paraffin)

Dilution: WB 1:500-2000, IF 1:50-300, IHC 1:50-300



IHC (Immunohistochemistry-Paraffin)

Dilution: WB 1:500-2000, IF 1:50-300, IHC 1:50-300

ADDITIONAL INFORMATION

Related Product Information for anti-CASP8 antibody

Most upstream protease of the activation cascade of caspases responsible for the TNFRSF6/FAS mediated and TNFRSF1A induced cell death. Binding to the adapter molecule FADD recruits it to either receptor. The resulting aggregate called death- inducing signaling complex (DISC) performs CASP8 proteolytic activation. The active dimeric enzyme is then liberated from the DISC and free to activate downstream apoptotic proteases. Proteolytic fragments of the N-terminal propeptide (termed CAP3, CAP5 and CAP6) are likely retained in the DISC. Cleaves and activates CASP3, CASP4, CASP6, CASP7, CASP9 and CASP10. May participate in the GZMB apoptotic pathways. Cleaves ADPRT. Hydrolyzes the small-molecule substrate, Ac-Asp-Glu-Val-Asp-I-AMC. Likely target for the cowpox virus CRMA death inhibitory protein. Isoform 5, isoform 6, isoform 7 and isoform 8 lack the catalytic site and may interfere with the pro-apoptotic activity of the complex.

Thiol protease that plays a key role in programmed cell death by acting as a molecular switch for apoptosis, necroptosis and pyroptosis, and is required to prevent tissue damage during embryonic development and adulthood (By similarity). Initiator protease that induces extrinsic apoptosis by mediating cleavage and activation of effector caspases responsible for the TNFRSF6/FAS mediated and TNFRSF1A induced cell death (PubMed:23516580, PubMed:8681376, PubMed:8681377, PubMed:9006941, PubMed:9184224, PubMed:8962078). Cleaves and activates effector caspases CASP3, CASP4, CASP6, CASP7, CASP9 and CASP10 (PubMed:8962078, PubMed:9006941). Binding to the adapter molecule FADD recruits it to either receptor TNFRSF6/FAS mediated or TNFRSF1A (PubMed:8681376, PubMed:8681377). The resulting aggregate called death- inducing signaling complex (DISC) performs CASP8 proteolytic activation (PubMed:9184224). The active dimeric enzyme is then liberated from the DISC and free to activate downstream apoptotic proteases (PubMed:9184224). Proteolytic fragments of the N-terminal propeptide (termed CAP3, CAP5 and CAP6) are likely retained in the DISC (PubMed:9184224). In addition to extrinsic apoptosis, also acts as a negative regulator of necroptosis: acts by cleaving RIPK1 at 'Asp-324', which is crucial to inhibit RIPK1 kinase activity, limiting TNF-induced apoptosis, necroptosis and inflammatory response (PubMed:31827280, PubMed:31827281). Also able to initiate pyroptosis by mediating cleavage and activation of gasdermin-D (GSDMD): GSDMD cleavage promoting release of the N-terminal moiety (Gasdermin-D, N-terminal) that binds to membranes and forms pores, triggering pyroptosis (By similarity). Initiates pyroptosis following inactivation of MAP3K7/TAK1 (By similarity). Also acts as a regulator of innate immunity by mediating cleavage and inactivation of N4BP1 downstream of TLR3 or TLR4, thereby promoting cytokine production (By similarity). May participate in the Granzyme B (GZMB) cell death pathways (PubMed:8755496). Cleaves PARP1 (PubMed:8681376).

NCBI AND UNIPROT INFORMATION

NCBI GI #

122056474

NCBI GeneID	841
NCBI Accession #	NP_001073593.1
NCBI GenBank Nucleotide #	NM_001080124.1
UniProt Accession #	Q14790; O14676; Q14791; Q14792; Q14793; Q14794; Q14795; Q14796; Q15780; Q15806; Q53TT5; Q8TDI1
Molecular Weight	55,391 Da
NCBI Official Full Name	caspase-8 isoform C
NCBI Official Synonym Full Names	caspase 8, apoptosis-related cysteine peptidase
NCBI Official Symbol	CASP8
NCBI Official Synonym Symbols	CAP4; MACH; MCH5; FLICE; ALPS2B; Casp-8
NCBI Protein Information	caspase-8; FADD-like ICE; MACH-alpha-1/2/3 protein; apoptotic protease Mch-5; MACH-beta-1/2/3/4 protein; apoptotic cysteine protease; ICE-like apoptotic protease 5; MORT1-associated ced-3 homolog; FADD-homologous ICE/CED-3-like protease; caspase 8, apopto
UniProt Protein Name	Caspase-8
UniProt Gene Name	CASP8
UniProt Synonym Gene Names	MCH5; CASP-8; FLICE; MACH
UniProt Entry Name	CASP8_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/cleaved-caspase-8-antibody-western-blot-immunohistochemistry-immunofluorescence-elisa/28833>