

Rabbit ATG4A Polyclonal Antibody | anti-ATG4A antibody

ATG4A Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA288688	Rabbit	Human, mouse (Predicted Reactivity: Bovine)	Immunohistochemistry, ELISA, Western Blot

PRODUCT SYNONYMS


ATG4A; Polyclonal Antibody; ATG4A Antibody; Cysteine protease ATG4A; 3422-; AUT-like 2 cysteine endopeptidase; Autophagin-2; Autophagy-related cysteine endopeptidase 2; Autophagy-related protein 4 homolog A; hAPG4A; APG4A; AUTL2; anti-ATG4A antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, mouse (Predicted Reactivity: Bovine)
Clonality	Polyclonal
Isotype	Rabbit Ig
Specificity	This ATG4A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 363-392 amino acids from human ATG4A.
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Concentration	Vial Concentration: 2 (varies by lot)
Sequence Positions	363-392
Sequence Length	398
Applicable Applications for anti-ATG4A antibody	IHC (Immunohistochemistry), ELISA, WB (Western Blot)
Antigen Type	Synthetic Peptide
Antigen Source	HUMAN
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 6 months. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.




WB (Western Blot)
APG4A Antibody (I378) western blot analysis in mouse kidney tissue lysates (35ug/lane). This demonstrates the APG4A antibody detected the APG4A protein (arrow).



IHC (Immunohistochemistry)
Formalin-fixed and paraffin-embedded human skeletal muscle tissue reacted with Autophagy APG4A Antibody (C-term) , which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



IHC (Immunohistochemistry)
Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.



WB (Western Blot)
Western blot analysis of anti-APG4A Pab in Jurkat cell line lysates (35ug/lane). APG4A (arrow) was detected using the purified Pab.

ADDITIONAL INFORMATION

Related Product Information for anti-ATG4A antibody	Macroautophagy is the major inducible pathway for the general turnover of cytoplasmic constituents in eukaryotic cells, it is also responsible for the degradation of active cytoplasmic enzymes and organelles during nutrient starvation. Macroautophagy involves the formation of double-membrane bound autophagosomes which enclose the cytoplasmic constituent targeted for degradation in a membrane bound structure, which then fuse with the lysosome (or vacuole) releasing a single-membrane bound autophagic bodies which are then degraded within the lysosome (or vacuole). APG4A is a cysteine protease required for autophagy, which cleaves the C-terminal part of either MAP1LC3, GABARAPL2 or GABARAP, allowing the liberation of form I. A subpopulation of form I is subsequently converted to a smaller form (form II). Form II, with a revealed C-terminal glycine, is considered to be the phosphatidylethanolamine (PE)-conjugated form, and has the capacity for the binding to autophagosomes. Preferred substrate is GABARAPL2 followed by MAP1LC3A and GABARAP.
Product Categories/Family for anti-ATG4A antibody	Cancer; Cardiovascular; Cell Biology; Metabolism; Autophagy
References	Baehrecke EH. Nat Rev Mol Cell Biol. 6(6):505-10. (2005) Lum JJ, et al. Nat Rev Mol Cell Biol. 6(6):439-48. (2005) Greenberg JT. Dev Cell. 8(6):799-801. (2005) Levine B. Cell. 120(2):159-62. (2005) Shintani T and Klionsky DJ. Science. 306(5698):990-5. (2004)

NCBI AND UNIPROT INFORMATION

NCBI GI #	30795252
NCBI GeneID	115201
NCBI Accession #	NP_443168.2; NP_840054.1
NCBI GenBank Nucleotide #	NM_052936.3
UniProt Accession #	Q8WYN0; Q95534; Q5JYY9; Q5JYZ0; Q86VE5; Q96KQ0; Q96KQ1; A6NCH2; B2RAZ7; D3DUY0
Molecular Weight	45378
NCBI Official Full Name	cysteine protease ATG4A isoform a
NCBI Official Synonym Full Names	autophagy related 4A, cysteine peptidase
NCBI Official Symbol	ATG4A
NCBI Official Synonym Symbols	APG4A; AUTL2
NCBI Protein Information	cysteine protease ATG4A
UniProt Protein Name	Cysteine protease ATG4A
UniProt Gene Name	ATG4A
UniProt Synonym Gene Names	APG4A; AUTL2; hAPG4A
UniProt Entry Name	ATG4A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/atg4a-antibody-immunohistochemistry-western-blot-elisa/288688>