

Rabbit anti-Human FTO Polyclonal Antibody | anti-FTO antibody

FTO Antibody (N-term)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA288780	Rabbit	Human	Flow Cytometry, Functional Assay, Immunohistochemistry, ELISA, Western Blot


PRODUCT SYNONYMS

FTO; Polyclonal Antibody; FTO Antibody (N-term); Alpha-ketoglutarate-dependent dioxygenase FTO; 11411-; Fat mass and obesity-associated protein; KIAA1752; anti-FTO antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit Ig
Specificity	This FTO antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human FTO.
Purity/Purification	Peptide Affinity Purified Rabbit Polyclonal Antibody (Pab)
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Concentration	Vial Concentration: 0.5 (varies by lot)
Sequence Positions	1-30
Sequence Length	505
Applicable Applications for anti-FTO antibody	FCM/FACS (Flow Cytometry), IHC (Immunohistochemistry), ELISA, WB (Western Blot)
Antigen Type	Synthetic Peptide
Antigen Source	HUMAN
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 6 months. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.


IMAGES



FCM/FACS (Flow Cytometry)
Flow cytometric analysis of NCI-H292 cells using FTO Antibody (N-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



IHC (Immunohistochemistry)
Formalin-fixed and paraffin-embedded human brain tissue reacted with FTO Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



WB (Western Blot)
Western blot analysis of FTO Antibody (N-term) in NCI-H460 cell line lysates (35ug/lane). FTO (arrow) was detected using the purified Pab.

ADDITIONAL INFORMATION

Related Product Information for anti-FTO antibody	The precise function of FTO remains to be determined.
Product Categories/Family for anti-FTO antibody	Cardiovascular ; Metabolism ; Neuroscience
References	Scott,L.J., et.al., Science 316 (5829), 1341-1345 (2007)

NCBI AND UNIPROT INFORMATION

NCBI GI #	122937263
NCBI GeneID	79068
NCBI Accession #	NP_001073901.1
NCBI GenBank Nucleotide #	NM_001080432.2
UniProt Accession #	Q9C0B1 ; Q0P676 ; Q7Z785 ; A2RUH1 ; B2RNS0
Molecular Weight	58282
NCBI Official Full Name	alpha-ketoglutarate-dependent dioxygenase FTO

NCBI Official Synonym Full Names	fat mass and obesity associated
NCBI Official Symbol	FTO
NCBI Official Synonym Symbols	ALKBH9
NCBI Protein Information	alpha-ketoglutarate-dependent dioxygenase FTO
UniProt Protein Name	Alpha-ketoglutarate-dependent dioxygenase FTO
UniProt Gene Name	FTO
UniProt Synonym Gene Names	KIAA1752
UniProt Entry Name	FTO_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/fto-antibody-flow-cytometry-immunohistochemistry-western-blot-elisa/288780>