

Rabbit EphA7 Polyclonal Antibody | anti-EPHA7 antibody

EphA7 Antibody (N-term)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA288791	Rabbit	Human Predicted Reactivity: Mouse, Rat	Flow Cytometry, Functional Assay, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

EphA7; Polyclonal Antibody; EphA7 Antibody (N-term); Ephrin type-A receptor 7; EPH homology kinase 3; EHK-3; EPH-like kinase 11; EK11; hEK11; EPHA7; EHK3; HEK11; anti-EPHA7 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human Predicted Reactivity: Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	This EphA7 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 11-41 amino acids from the N-terminal region of human EphA7.
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Sequence Positions	11-41
Sequence Length	998
Applicable Applications for anti-EPHA7 antibody	FCM/FACS (Flow Cytometry), IHC (Immunohistochemistry), ELISA
Function	Receptor tyrosine kinase which binds promiscuously GPI- anchored ephrin-A family ligands residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. Among GPI-anchored ephrin-A ligands, EFNA5 is a cognate/functional ligand for EPHA7 and their interaction regulates brain development modulating cell-cell adhesion and repulsion. Has a repellent activity on axons and is for instance involved in the guidance of corticothalamic axons and in the proper topographic mapping of retinal axons to the colliculus. May also regulate brain development through a caspase(CASP3)-dependent proapoptotic activity. Forward signaling may result in activation of components of the ERK signaling pathway including MAP2K1, MAP2K2, MAPK1 AND MAPK3 which are phosphorylated upon activation of EPHA7.
Cellular Location	Cell membrane; Single-pass type I membrane protein
Tissue Location	Widely expressed.
Preparation and Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

IMAGES



IHC (Immunohiostchemistry)
Formalin-fixed and paraffin-embedded human brain tissue reacted with EphA7 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



FCM/FACS (Flow Cytometry)
Flow cytometric analysis of A375 cells using EphA7 Antibody (N-term) (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

ADDITIONAL INFORMATION

Related Product Information for anti-EPHA7 antibody	Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the g phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains. The tyrosine kinase (TK) group is mainly involved in the regulation of cell-cell interactions such as differentiation, adhesion, motility and death. There are currently about 90 TK genes sequenced, 58 are of receptor protein TK (e.g. EGFR, EPH, FGFR, PDGFR, TRK, and VEGFR families), and 32 of cytosolic TK (e.g. ABL, FAK, JAK, and SRC families).
Product Categories/Family for anti-EPHA7 antibody	Cardiovascular; Neuroscience
References	Wilkinson, D.G., Nat Rev Neurosci 2(3):155-164 (2001). Xu, Q., et al., Philos. Trans. R. Soc. Lond., B, Biol. Sci. 355(1399):993-1002 (2000). Holder, N., et al., Development 126(10):2033-2044 (1999). Zhou, R., Pharmacol. Ther. 77(3):151-181 (1998). Fox, G.M., et al., Oncogene 10(5):897-905 (1995).

NCBI AND UNIPROT INFORMATION

NCBI GI #	4758282
NCBI GeneID	2045
NCBI Accession #	Q15375
NCBI GenBank Nucleotide #	NM_004440.3
NCBI Official Full Name	ephrin type-A receptor 7 isoform 1
NCBI Official Synonym Full Names	EPH receptor A7
NCBI Official Symbol	EPHA7
NCBI Official Synonym Symbols	EHK3; EK11; EHK-3; HEK11
NCBI Protein Information	ephrin type-A receptor 7
UniProt Protein Name	Ephrin type-A receptor 7
UniProt Gene Name	EPHA7
UniProt Synonym Gene Names	EHK3; HEK11; EHK-3; EK11; hEK11

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/epha7-antibody-flow-cytometry-immunohistochemistry-elisa/288791>