

Rabbit anti-Human Bad Polyclonal Antibody | anti-BAD antibody

Bad Antibody (Center)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA289024	Rabbit	Human	Immunohistochemistry, ELISA, Western Blot


PRODUCT SYNONYMS

Bad; Polyclonal Antibody; Bad Antibody (Center); Bcl2-associated agonist of cell death; BAD; Bcl-2-binding component 6; Bcl-2-like protein 8; Bcl2-L-8; Bcl-xL/Bcl-2-associated death promoter; Bcl2 antagonist of cell death; BBC6; BCL2L8; anti-BAD antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	This Bad antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 53-81 amino acids from the Central region of human Bad.
Purity/Purification	Peptide Affinity Purified Rabbit Polyclonal Antibody (Pab)
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Concentration	Vial Concentration: 0.38 (varies by lot)
Sequence Positions	53-81
Sequence Length	168
Applicable Applications for anti-BAD antibody	IHC (Immunohistochemistry), ELISA, WB (Western Blot)
Antigen Type	Synthetic Peptide
Antigen Source	HUMAN
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 6 months. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



IHC (Immunohiostchemistry)
Formalin-fixed and paraffin-embedded human lung carcinoma tissue reacted with Bad antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



WB (Western Blot)
Western blot analysis of anti-Bad Antibody (Center) in HL60 cell line lysates (35ug/lane). Bad(arrow) was detected using the purified Pab.

ADDITIONAL INFORMATION

Related Product Information for anti-BAD antibody	Bad is a member of the BCL-2 family. BCL-2 family members are known to be regulators of programmed cell death. This protein positively regulates cell apoptosis by forming heterodimers with BCL-xL and BCL-2, and reversing their death repressor activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be involved in the regulation of this protein.
Product Categories/Family for anti-BAD antibody	Cancer; Cell Biology; Metabolism; Signal Transduction; Amyotrophic Lateral Sclerosis (ALS)
References	Lee, J.W., et al., Carcinogenesis 25(8):1371-1376 (2004). Zhang, B., et al., Mol. Cell. Biol. 24(14):6205-6214 (2004). Ong, C.S., et al., Oncol. Rep. 11(3):727-733 (2004). Yan, B., et al., J. Biol. Chem. 278(46):45358-45367 (2003). Taghiyev, A.F., et al., Mol. Cancer Res. 1(7):500-507 (2003).

NCBI AND UNIPROT INFORMATION

NCBI GI #	10835069
NCBI GeneID	572
NCBI Accession #	NP_004313.1; NP_116784.1
NCBI GenBank Nucleotide #	NM_004322.3
UniProt Accession #	Q92934; O14803; Q6FH21
Molecular Weight	18392
NCBI Official Full Name	bcl2 antagonist of cell death
NCBI Official Synonym Full Names	BCL2-associated agonist of cell death
NCBI Official Symbol	BAD
NCBI Official Synonym Symbols	BBC2; BCL2L8
NCBI Protein Information	bcl2-associated agonist of cell death
UniProt Protein Name	Bcl2-associated agonist of cell death
UniProt Gene Name	BAD

UniProt Synonym Gene Names	BBC6; BCL2L8; BAD; Bcl2-L-8
UniProt Entry Name	BAD_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/bad-antibody-immunohistochemistry-western-blot-elisa/289024>