

Rabbit PCNA Polyclonal Antibody | anti-PCNA antibody

PCNA Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA290119	Rabbit	Human, mouse, rat	Immunohistochemistry, ELISA, Western Blot


PRODUCT SYNONYMS

PCNA; Polyclonal Antibody; PCNA Antibody; Proliferating cell nuclear antigen; Cyclin; anti-PCNA antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, mouse, rat
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Form/Format	Rabbit IgG in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Sequence Positions	80-115
Sequence Length	261
Applicable Applications for anti-PCNA antibody	IHC (Immunohistochemistry), ELISA, WB (Western Blot)
Antigen Type	Synthetic Peptide
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 6 months. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using PCNA antibody.



WB (Western Blot)

Western blot analysis of extracts from Hela cells (Lane 2), colo cells(Lane 3), HepG2 cells (Lane 4) and 293 cells (Lane 5) using PCNA Antibody. The lane on the left is treated with synthesized peptide.



WB (Western Blot)

Western blot analysis of lysates from HeLa, MCF-7 cell line (from left to right), using PCNA Antibody (C0298). C0298 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35 µg per lane.

ADDITIONAL INFORMATION

Related Product Information for anti-PCNA antibody	Auxiliary protein of DNA polymerase delta and is involved in the control of eukaryotic DNA replication by increasing the polymerase's processibility during elongation of the leading strand. Induces a robust stimulatory effect on the 3'-5' exonuclease and 3'-phosphodiesterase, but not apurinic- apyrimidinic (AP) endonuclease, APEX2 activities. Has to be loaded onto DNA in order to be able to stimulate APEX2. Plays a key role in DNA damage response (DDR) by being conveniently positioned at the replication fork to coordinate DNA replication with DNA repair and DNA damage tolerance pathways. Acts as a loading platform to recruit DDR proteins that allow completion of DNA replication after DNA damage and promote postreplication repair: Monoubiquitinated PCNA leads to recruitment of translesion (TLS) polymerases, while 'Lys-63'-linked polyubiquitination of PCNA is involved in error-free pathway and employs recombination mechanisms to synthesize across the lesion.
Product Categories/Family for anti-PCNA antibody	Cancer ; Cell biology
References	Almendral J.M., et al. Proc. Natl. Acad. Sci. U.S.A. 84:1575-1579(1987). Travali S., et al. J. Biol. Chem. 264:7466-7472(1989). Ota T., et al. Nat. Genet. 36:40-45(2004). Deloukas P., et al. Nature 414:865-871(2001). Mural R.J., et al. Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4505641
NCBI GeneID	5111
NCBI Accession #	NP_002583.1
NCBI GenBank Nucleotide #	NM_002592.2
UniProt Accession #	P12004 ; B2R897 ; D3DW02
Molecular Weight	28769
NCBI Official Full Name	proliferating cell nuclear antigen
NCBI Official Synonym Full Names	proliferating cell nuclear antigen
NCBI Official Symbol	PCNA
NCBI Official Synonym Symbols	ATLD2
NCBI Protein Information	proliferating cell nuclear antigen
UniProt Protein Name	Proliferating cell nuclear antigen

UniProt Gene Name	PCNA
UniProt Synonym Gene Names	PCNA
UniProt Entry Name	PCNA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/pcna-antibody-immunohistochemistry-western-blot-elisa/290119>