

Rabbit EPHA2 Polyclonal Antibody | anti-EPHA2 antibody

EPHA2 Antibody (Center)


CATALOG # AAA290358	HOST Rabbit	REACTIVITY Human (Predicted Reactivity: Monkey)	APPLICATIONS ELISA, Western Blot
-------------------------------	-----------------------	---	--

PRODUCT SYNONYMS

EPHA2; Polyclonal Antibody; EPHA2 Antibody (Center); Ephrin type-A receptor 2; Epithelial cell kinase; Tyrosine-protein kinase receptor ECK; ECK; anti-EPHA2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human (Predicted Reactivity: Monkey)
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	This EPHA2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 518-552 amino acids from the Central region of human EPHA2.
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Form/Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Sequence Length	497
Applicable Applications for anti-EPHA2 antibody	ELISA, WB (Western Blot)
Function	Receptor tyrosine kinase which binds promiscuously membrane- bound ephrin-A family ligands residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. Activated by the ligand ephrin- A1/EFNA1 regulates migration, integrin-mediated adhesion, proliferation and differentiation of cells. Regulates cell adhesion and differentiation through DSG1/desmoglein-1 and inhibition of the ERK1/ERK2 (MAPK3/MAPK1, respectively) signaling pathway. May also participate in UV radiation-induced apoptosis and have a ligand-independent stimulatory effect on chemotactic cell migration. During development, may function in distinctive aspects of pattern formation and subsequently in development of several fetal tissues. Involved for instance in angiogenesis, in early hindbrain development and epithelial proliferation and branching morphogenesis during mammary gland development. Engaged by the ligand ephrin-A5/EFNA5 may regulate lens fiber cells shape and interactions and be important for lens transparency development and maintenance. With ephrin-A2/EFNA2 may play a role in bone remodeling through regulation of osteoclastogenesis and osteoblastogenesis.
Cellular Location	Cell membrane; Single-pass type I membrane protein. Cell projection, ruffle membrane; Single-pass type I membrane protein. Cell projection, lamellipodium membrane; Single-pass type I membrane protein. Cell junction, focal adhesion. Note=Present at regions of cell-cell contacts but also at the leading edge of migrating cells (PubMed:19573808, PubMed:20861311). Relocates from the plasma membrane to the cytoplasmic and perinuclear regions in cancer cells (PubMed:18794797).
Tissue Location	Expressed in brain and glioma tissue and glioma cell lines (at protein level). Expressed most highly in tissues that contain a high proportion of epithelial cells, e.g. skin, intestine, lung, and ovary.
Preparation and Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.



WB (Western Blot)
Western blot analysis of lysates from A431, U-87 MG cell line (from left to right), using EPHA2 Antibody (Center). AAA290358 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

ADDITIONAL INFORMATION

Related Product Information for anti-EPHA2 antibody	Receptor tyrosine kinase which binds promiscuously membrane-bound ephrin-A family ligands residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. Activated by the ligand ephrin-A1/EFNA1 regulates migration, integrin-mediated adhesion, proliferation and differentiation of cells. Regulates cell adhesion and differentiation through DSG1/desmoglein-1 and inhibition of the ERK1/ERK2 (MAPK3/MAPK1, respectively) signaling pathway. May also participate in UV radiation-induced apoptosis and have a ligand- independent stimulatory effect on chemotactic cell migration. During development, may function in distinctive aspects of pattern formation and subsequently in development of several fetal tissues. Involved for instance in angiogenesis, in early hindbrain development and epithelial proliferation and branching morphogenesis during mammary gland development. Engaged by the ligand ephrin-A5/EFNA5 may regulate lens fiber cells shape and interactions and be important for lens transparency development and maintenance. With ephrin-A2/EFNA2 may play a role in bone remodeling through regulation of osteoclastogenesis and osteoblastogenesis.
Product Categories/Family for anti-EPHA2 antibody	Cardiovascular; Neuroscience; Signal Transduction; Stem Cells
References	Lindberg R.A.,et al.Mol. Cell. Biol. 10:6316-6324(1990). Gregory S.G.,et al.Nature 441:315-321(2006). Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases. Miao H.,et al.Nat. Cell Biol. 2:62-69(2000). Zelinski D.P.,et al.Cancer Res. 61:2301-2306(2001).

NCBI AND UNIPROT INFORMATION

NCBI GI #	32967311
NCBI GeneID	1969
NCBI Accession #	NP_004422.2
NCBI GenBank Nucleotide #	NM_004431.3
UniProt Accession #	P29317; Q8N3Z2; B5A968
Molecular Weight	108266
NCBI Official Full Name	ephrin type-A receptor 2
NCBI Official Synonym Full Names	EPH receptor A2

NCBI Official Symbol	EPHA2
NCBI Official Synonym Symbols	ECK; CTPA; ARCC2; CTPP1; CTRCT6
NCBI Protein Information	ephrin type-A receptor 2
UniProt Protein Name	Ephrin type-A receptor 2
UniProt Gene Name	EPHA2
UniProt Synonym Gene Names	ECK
UniProt Entry Name	EPHA2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/epha2-antibody-western-blot-elisa/290358>