

Rabbit anti-Rat, Dog FCGBP Polyclonal Antibody | anti-FCGBP antibody

FCGBP Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291455	Rabbit	Rat, Dog	Immunofluorescence, Immunohistochemistry, Immunohistochemistry


PRODUCT SYNONYMS

FCGBP; Polyclonal Antibody; FCGBP Polyclonal Antibody; IgGfC-binding protein; Fcgamma-binding protein antigen; FcgammaBP; anti-FCGBP antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat, Dog
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-FCGBP antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES





IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (mouse intestine); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (FCGBP) Polyclonal Antibody, Unconjugated (bs-13168R) at 1:200 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Dilution: Tissue/cell: human gastric carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37 degree C for 20 min; Incubation: Anti-FCGBP Polyclonal Antibody, Unconjugated (bs-13168R) 1:200, overnight at 4 degree C, followed by conjugation to the secondary antibody (SP-0023) and DAB (C-0010) staining



IHC (Immunohistochemistry)

Positive sample? Paraformaldehyde-fixed, paraffin embedded (rat intestine); Negative sample? Paraformaldehyde-fixed, paraffin embedded (rat lung); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (FCGBP) Polyclonal Antibody, Unconjugated (bs-13168R) at 1:200 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

IHC (Immunohistochemistry)

Tissue/cell: human gastric carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37 degree C for 20 min; Incubation: Anti-FCGBP Polyclonal Antibody, Unconjugated (bs-13168R) 1:200, overnight at 4 degree C, followed by conjugation to the secondary antibody (SP-0023) and DAB (C-0010) staining

NCBI AND UNIPROT INFORMATION

NCBI GI #	119577330
NCBI GeneID	8857
UniProt Accession #	Q9Y6R7; O95784
Molecular Weight	572,017 Da
NCBI Official Full Name	Fc fragment of IgG binding protein
NCBI Official Synonym Full Names	Fc fragment of IgG binding protein
NCBI Official Symbol	FCGBP
NCBI Official Synonym Symbols	FC(GAMMA)BP
NCBI Protein Information	IgGFc-binding protein
UniProt Protein Name	IgGFc-binding protein
UniProt Gene Name	FCGBP
UniProt Synonym Gene Names	FcgammaBP
UniProt Entry Name	FCGBP_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/fcgbp-antibody-immunofluorescence-immunohistochemistry-immunohistochemistry-elisa/291455>