

Rabbit Keap1 Polyclonal Antibody | anti-Keap1 antibody

Keap1 Polyclonal Antibody

CATALOG # AAA29182	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Western Blot, Immunohistochemistry
------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

Keap1; Polyclonal Antibody; Keap1 Polyclonal Antibody; KEAP1; INRF2; KIAA0132; KLHL19; Kelch-like ECH-associated protein 1; Cytosolic inhibitor of Nrf2; INrf2; Kelch-like protein 19; anti-Keap1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-Keap1 antibody	WB (Western Blot), IHC (Immunohistochemistry)
Notes	ECO:0000303 PubMed:14585973, ECO:0000312 HGNC: HGNC:23177
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.




WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-Keap1 antibody

Acts as a substrate adapter protein for the E3 ubiquitin ligase complex formed by CUL3 and RBX1 and targets NFE2L2/NRF2 for ubiquitination and degradation by the proteasome, thus resulting in the suppression of its transcriptional activity and the repression of antioxidant response element-mediated detoxifying enzyme gene expression. Retains NFE2L2/NRF2 and may also retain BPTF in the cytosol. Targets PGAM5 for ubiquitination and degradation by the proteasome.

NCBI AND UNIPROT INFORMATION

NCBI GI #	146345444
NCBI GeneID	9817
UniProt Accession #	Q14145; Q6LEP0; Q8WTX1; Q9BPY9; B3KPD5
Molecular Weight	624
NCBI Official Full Name	Kelch-like ECH-associated protein 1
NCBI Official Synonym Full Names	kelch-like ECH-associated protein 1
NCBI Official Symbol	KEAP1
NCBI Official Synonym Symbols	INrf2; KLHL19
NCBI Protein Information	kelch-like ECH-associated protein 1; kelch-like protein 19; cytosolic inhibitor of Nrf2; kelch-like family member 19
UniProt Protein Name	Kelch-like ECH-associated protein 1
UniProt Gene Name	KEAP1
UniProt Synonym Gene Names	INRF2; KIAA0132; KLHL19; INrf2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/keap1-antibody-western-blot-immunohistochemistry-elisa/29182>