

Rabbit PSMA8 Polyclonal Antibody | anti-PSMA8 antibody

PSMA8 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA291844	Rabbit	Rat, Pig, Cynomolgus Monkey, Dog, Cow	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

PSMA8; Polyclonal Antibody; PSMA8 Polyclonal Antibody; Proteasome subunit alpha-type 8; Proteasome alpha 4 subunit; Proteasome subunit alpha-type 7-like; PSMA7L; anti-PSMA8 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat, Pig, Cynomolgus Monkey, Dog, Cow
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-PSMA8 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (PSMA8) Polyclonal Antibody, Unconjugated (bs-19461R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (Mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (PSMA8) Polyclonal Antibody, Unconjugated (bs-19461R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



WB (Western Blot)
 Sample:TM4(Mouse) Cell Lysate at 30 ugPrimary: Anti-PSMA8 (bs-19461R) at 1/300 dilutionSecondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilutionPredicted band size: 29 kD
 Observed band size: 29 kD

NCBI AND UNIPROT INFORMATION

NCBI GI #	68303563
NCBI GeneID	143471
NCBI Accession #	NP_001020267.1
NCBI GenBank Nucleotide #	NM_001025096.1
UniProt Accession #	Q8TAA3; Q8IVP4; Q8TA98; Q8TAA2; B0YJ75
Molecular Weight	27,843 Da
NCBI Official Full Name	proteasome subunit alpha type-7-like isoform 2
NCBI Official Synonym Full Names	proteasome (prosome, macropain) subunit, alpha type, 8
NCBI Official Symbol	PSMA8
NCBI Official Synonym Symbols	PSMA7L
NCBI Protein Information	proteasome subunit alpha type-7-like
UniProt Protein Name	Proteasome subunit alpha type-7-like
UniProt Gene Name	PSMA8
UniProt Synonym Gene Names	PSMA7L
UniProt Entry Name	PSA7L_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://www.aaabiotech.com/polyclonal/psma8-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/291844>