

Rabbit anti-Human CD2 Polyclonal Antibody | anti-CD2 antibody

CD2 Polyclonal Antibody

CATALOG # AAA29190	HOST Rabbit	REACTIVITY Human	APPLICATIONS Western Blot, Immunohistochemistry
------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

CD2; Polyclonal Antibody; CD2 Polyclonal Antibody; SRBC; T-cell surface antigen CD2; Erythrocyte receptor; LFA-2; LFA-3 receptor; Rosette receptor; T-cell surface antigen T11/Leu-5; anti-CD2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Applicable Applications for anti-CD2 antibody	WB (Western Blot), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES





IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.


Application Data

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-CD2 antibody

CD2 interacts with lymphocyte function-associated antigen CD58 (LFA-3) and CD48/BCM1 to mediate adhesion between T- cells and other cell types. CD2 is implicated in the triggering of T-cells, the cytoplasmic domain is implicated in the signaling function.

NCBI AND UNIPROT INFORMATION

NCBI GI #	156071472
NCBI GeneID	914
NCBI Accession #	NP_001758.2
NCBI GenBank Nucleotide #	NM_001767.3
UniProt Accession #	P06729; Q96TE5
Molecular Weight	39,448 Da
NCBI Official Full Name	T-cell surface antigen CD2
NCBI Official Synonym Full Names	CD2 molecule
NCBI Official Symbol	CD2
NCBI Official Synonym Symbols	T11; SRBC; LFA-2
NCBI Protein Information	T-cell surface antigen CD2; CD2 antigen (p50), sheep red blood cell receptor; LFA-3 receptor; T-cell surface antigen T11/Leu-5; erythrocyte receptor; lymphocyte-function antigen-2; rosette receptor
UniProt Protein Name	T-cell surface antigen CD2

UniProt Gene Name	CD2
UniProt Synonym Gene Names	SRBC
UniProt Entry Name	CD2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/cd2-antibody-western-blot-immunohistochemistry-elisa/29190>