

Rabbit Integrin alpha4 Polyclonal Antibody | anti-ITGA4 antibody

Integrin alpha4 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA29192	Rabbit	Human, Mouse, Rat	Western Blot, Immunohistochemistry


PRODUCT SYNONYMS

Integrin alpha4; Polyclonal Antibody; Integrin alpha4 Polyclonal Antibody; ITGA4; CD49D; Integrin alpha-4; CD49 antigen-like family member D; Integrin alpha-IV; VLA-4 subunit alpha; CD49d; anti-ITGA4 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-ITGA4 antibody	WB (Western Blot), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-ITGA4 antibody

Integrins alpha-4/beta-1 (VLA-4) and alpha-4/beta-7 are receptors for fibronectin. They recognize one or more domains within the alternatively spliced CS-1 and CS-5 regions of fibronectin. They are also receptors for VCAM1. Integrin alpha- 4/beta-1 recognizes the sequence Q-I-D-S in VCAM1. Integrin alpha- 4/beta-7 is also a receptor for MADCAM1. It recognizes the sequence L-D-T in MADCAM1. On activated endothelial cells integrin VLA-4 triggers homotypic aggregation for most VLA-4-positive leukocyte cell lines. It may also participate in cytolytic T-cell interactions with target cells. ITGA4: ITGB1 binds to fractalkine (CX3CL1) and may act as its coreceptor in CX3CR1-dependent fractalkine signaling (PubMed:25398877).

NCBI AND UNIPROT INFORMATION

NCBI GI #	311033436
NCBI GeneID	3676
UniProt Accession #	P13612; Q7Z4L6; D3DPG4; AAB25486
Molecular Weight	1032
NCBI Official Full Name	Integrin alpha-4
NCBI Official Synonym Full Names	integrin, alpha 4 (antigen CD49D, alpha 4 subunit of VLA-4 receptor)
NCBI Official Symbol	ITGA4
NCBI Official Synonym Symbols	IA4; CD49D
NCBI Protein Information	integrin alpha-4; 269C wild type; integrin alpha 4; integrin alpha-IV; VLA-4 subunit alpha; integrin alpha-4 subunit; CD49 antigen-like family member D; antigen CD49D, alpha-4 subunit of VLA-4 receptor; very late activation protein 4 receptor, alpha 4 sub
UniProt Protein Name	Integrin alpha-4

UniProt Gene Name	ITGA4
UniProt Synonym Gene Names	CD49D
UniProt Entry Name	ITA4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/integrin-alpha4-antibody-western-blot-immunohistochemistry-elisa/29192>