

Rabbit anti-Human CD79b Polyclonal Antibody | anti-CD79b antibody

CD79b Polyclonal Antibody

CATALOG # AAA29197	HOST Rabbit	REACTIVITY Human	APPLICATIONS Western Blot, Immunohistochemistry
------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

CD79b; Polyclonal Antibody; CD79b Polyclonal Antibody; CD79B; B29; IGB; B-cell antigen receptor complex-associated protein beta chain; B-cell-specific glycoprotein B29; Ig-beta; Immunoglobulin-associated B29 protein; anti-CD79b antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Applicable Applications for anti-CD79b antibody	WB (Western Blot), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES





IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.


Application Data

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry-Paraffin)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-CD79b antibody

Required in cooperation with CD79A for initiation of the signal transduction cascade activated by the B-cell antigen receptor complex (BCR) which leads to internalization of the complex, trafficking to late endosomes and antigen presentation. Enhances phosphorylation of CD79A, possibly by recruiting kinases which phosphorylate CD79A or by recruiting proteins which bind to CD79A and protect it from dephosphorylation.

NCBI AND UNIPROT INFORMATION

NCBI GI #	728994
NCBI GeneID	974
UniProt Accession #	P40259; Q53FS2; Q9BU06
Molecular Weight	229
NCBI Official Full Name	B-cell antigen receptor complex-associated protein beta chain
NCBI Official Synonym Full Names	CD79b molecule, immunoglobulin-associated beta
NCBI Official Symbol	CD79B
NCBI Official Synonym Symbols	B29; IGB; AGM6
NCBI Protein Information	B-cell antigen receptor complex-associated protein beta chain; Ig-beta; B-cell-specific glycoprotein B29; immunoglobulin-associated B29 protein; CD79b antigen (immunoglobulin-associated beta)
UniProt Protein Name	B-cell antigen receptor complex-associated protein beta chain
UniProt Gene Name	CD79B

UniProt Synonym Gene Names	B29; IGB
UniProt Entry Name	CD79B_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/cd79b-antibody-western-blot-immunohistochemistry-elisa/29197>