

Rabbit anti-Rat NPD014 Polyclonal Antibody | anti-NPD014 antibody

NPD014 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA292042	Rabbit	Rat	Immunofluorescence, Immunohistochemistry, Immunohistochemistry


PRODUCT SYNONYMS

NPD014; Polyclonal Antibody; NPD014 Polyclonal Antibody; Arginine/serine-rich protein 1; RSRP1; C1orf63; anti-NPD014 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-NPD014 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES





IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (Rat stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (NPD014) Polyclonal Antibody, Unconjugated (bs-6441R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Dilution: Paraformaldehyde-fixed, paraffin embedded (Rat bladder); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (NPD014) Polyclonal Antibody, Unconjugated (bs-6441R) at 1:400 overnight at 4 degree C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



IHC (Immunohistochemistry)

Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37 degree C for 20 min;Incubation: Anti-NPD014 Polyclonal Antibody, Unconjugated(bs-6441R) 1:200, overnight at 4 degree C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

IHC (Immunohistochemistry)

Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37 degree C for 20 min;Incubation: Anti-NPD014 Polyclonal Antibody, Unconjugated(bs-6441R) 1:200, overnight at 4 degree C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

NCBI AND UNIPROT INFORMATION

NCBI GI #	46309849
NCBI GeneID	57035
NCBI Accession #	NP_064713.3
NCBI GenBank Nucleotide #	NM_020317.3
UniProt Accession #	Q9BUV0; Q49AA4; Q5TH71; Q9GZP6; A8K917
Molecular Weight	26,605 Da
NCBI Official Full Name	arginine/serine-rich protein 1
NCBI Official Synonym Full Names	arginine/serine-rich protein 1
NCBI Official Symbol	RSRP1
NCBI Official Synonym Symbols	NPD014; C1orf63; DJ465N24.2.1
NCBI Protein Information	arginine/serine-rich protein 1; UPF0471 protein C1orf63
UniProt Protein Name	Arginine/serine-rich protein 1
UniProt Gene Name	RSRP1
UniProt Synonym Gene Names	C1orf63
UniProt Entry Name	RSRP1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/npd014-antibody-immunofluorescence-immunohistochemistry-immunohistochemistry-elisa/292042>