

Rabbit anti-Rat, Cow KRCC1 Polyclonal Antibody | anti-KRCC1 antibody

KRCC1 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA292201	Rabbit	Rat, Cow	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

KRCC1; Polyclonal Antibody; KRCC1 Polyclonal Antibody; Lysine-rich coiled-coil protein 1; Cryptogenic hepatitis-binding protein 2; CHBP2; anti-KRCC1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Rat, Cow
Clonality	Polyclonal
Purity/Purification	Purified Rabbit Polyclonal Antibody (Pab)
Applicable Applications for anti-KRCC1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (KRCC1) Polyclonal Antibody, Unconjugated (bs-9560R) at 1:400 overnight at 4 degree C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.




IHC (Immunohistochemistry)

Dilution: Paraformaldehyde-fixed, paraffin embedded (Mouse Kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (KRCC1) Polyclonal Antibody, Unconjugated (bs9560R) at 1:400 overnight at 4 degree C, followed by a conjugated secondary antibody (sp-0023) for 20 minutes and DAB staining.



IHC (Immunohistochemistry)

Paraformaldehyde-fixed, paraffin embedded (Mouse Brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 degree C for 30min; Antibody incubation with (KRCC1) Polyclonal Antibody, Unconjugated (bs9560R) at 1:400 overnight at 4 degree C, followed by a conjugated secondary antibody (sp-0023) for 20 minutes and DAB staining.



WB (Western Blot)

Sample: Kidney (Mouse) Lysate at 40 ug Cerebrum (Mouse) Lysate at 40 ug
 Primary: Anti-KRCC1 (bs-9560R) at 1/300 dilution
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
 Predicted band size: 31 kDa
 Observed band size: 31 kDa

NCBI AND UNIPROT INFORMATION

NCBI GI #	7706156
NCBI GeneID	51315
NCBI Accession #	NP_057702.1
NCBI GenBank Nucleotide #	NM_016618.1
UniProt Accession #	Q9NPI7; Q3B7J7
Molecular Weight	30,980 Da
NCBI Official Full Name	lysine-rich coiled-coil protein 1
NCBI Official Synonym Full Names	lysine-rich coiled-coil 1
NCBI Official Symbol	KRCC1
NCBI Official Synonym Symbols	CHBP2
NCBI Protein Information	lysine-rich coiled-coil protein 1; cryptogenic hepatitis binding protein; cryptogenic hepatitis-binding protein 2
UniProt Protein Name	Lysine-rich coiled-coil protein 1
UniProt Gene Name	KRCC1
UniProt Synonym Gene Names	CHBP2; BM-003; BM-044
UniProt Entry Name	KRCC1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/krcc1-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-immunohistochemistry-western-blot-elisa/292201>