

Rabbit CRM1 Polyclonal Antibody | anti-CRM1 antibody

CRM1 (Acetyl Lys568) Polyclonal Antibody

CATALOG # AAA293497	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Western Blot
-------------------------------	-----------------------	--	-------------------------------------


PRODUCT SYNONYMS

CRM1; Polyclonal Antibody; CRM1 (Acetyl Lys568) Polyclonal Antibody; XPO1; Exportin-1; Exp1; Chromosome region maintenance 1 protein homolog; anti-CRM1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Form/Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Applicable Applications for anti-CRM1 antibody	WB (Western Blot)
Cellular Location	Cytoplasm. Nucleus, nucleoplasm. Nucleus, Cajal body. Nucleus, nucleolus. Note=Located in the nucleoplasm, Cajal bodies and nucleoli. Shuttles between the nucleus/nucleolus and the cytoplasm
Tissue Location	Expressed in heart, brain, placenta, lung, liver, skeletal muscle, pancreas, spleen, thymus, prostate, testis, ovary, small intestine, colon and peripheral blood leukocytes. Not expressed in the kidney.
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-CRM1 antibody	Mediates the nuclear export of cellular proteins (cargos) bearing a leucine-rich nuclear export signal (NES) and of RNAs. In the nucleus, in association with RANBP3, binds cooperatively to the NES on its target protein and to the GTPase RAN in its active GTP-bound form (Ran-GTP). Docking of this complex to the nuclear pore complex (NPC) is mediated through binding to nucleoporins. Upon transit of a nuclear export complex into the cytoplasm, disassembling of the complex and hydrolysis of Ran-GTP to Ran-GDP (induced by RANBP1 and RANGAP1, respectively) cause release of the cargo from the export receptor. The directionality of nuclear export is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the cytoplasm and nucleus. Involved in U3 snoRNA transport from Cajal bodies to nucleoli. Binds to late precursor U3 snoRNA bearing a TMG cap.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	68052989
NCBI GeneID	7514
UniProt Accession #	O14980; Q63HP8; Q68CP3; Q99433; A6NL14; A8K1K5; D6W5E2
Molecular Weight	1071
NCBI Official Full Name	Exportin-1
NCBI Official Synonym Full Names	exportin 1
NCBI Official Symbol	XPO1
NCBI Official Synonym Symbols	emb; CRM1; exp1
NCBI Protein Information	exportin-1; exportin 1 (CRM1 homolog, yeast); chromosome region maintenance 1 homolog; chromosome region maintenance 1 protein homolog; exportin-1 (required for chromosome region maintenance)
UniProt Protein Name	Exportin-1
UniProt Gene Name	XPO1
UniProt Synonym Gene Names	CRM1; Exp1
UniProt Entry Name	XPO1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.