

Rabbit PYK2 Polyclonal Antibody | anti-PYK2 antibody

PYK2 (phospho Tyr881) Polyclonal Antibody

CATALOG # AAA294036	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

PYK2; Polyclonal Antibody; PYK2 (phospho Tyr881) Polyclonal Antibody; PTK2B; FAK2; RAFTK; Protein-tyrosine kinase 2-beta; Calcium-dependent tyrosine kinase; CADTK; Calcium-regulated non-receptor proline-rich tyrosine kinase; Cell adhesion kinase beta; CAK-beta; CAKB; Focal adhesion kinase 2; FADK 2; Pro; anti-PYK2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-PYK2 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.


IMAGES



IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000.

Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000.

Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-PYK2 antibody	Non-receptor protein-tyrosine kinase that regulates reorganization of the actin cytoskeleton, cell polarization, cell migration, adhesion, spreading and bone remodeling. Plays a role in the regulation of the humoral immune response, and is required for normal levels of marginal B-cells in the spleen and normal migration of splenic B-cells. Required for normal macrophage polarization and migration towards sites of inflammation. Regulates cytoskeleton rearrangement and cell spreading in T- cells, and contributes to the regulation of T-cell responses. Promotes osteoclastic bone resorption; this requires both PTK2B/PYK2 and SRC. May inhibit differentiation and activity of osteoprogenitor cells. Functions in signaling downstream of integrin and collagen receptors, immune receptors, G-protein coupled receptors (GPCR), cytokine, chemokine and growth factor receptors, and mediates responses to cellular stress. Forms multisubunit signaling complexes with SRC and SRC family members upon activation; this leads to the phosphorylation of additional tyrosine residues, creating binding sites for scaffold proteins, effectors and substrates. Regulates numerous signaling pathways. Promotes activation of phosphatidylinositol 3-kinase and of the AKT1 signaling cascade. Promotes activation of NOS3. Regulates production of the cellular messenger cGMP. Promotes activation of the MAP kinase signaling cascade, including activation of MAPK1/ERK2, MAPK3/ERK1 and MAPK8/JNK1. Promotes activation of Rho family GTPases, such as RHOA and RAC1. Recruits the ubiquitin ligase MDM2 to P53/TP53 in the nucleus, and thereby regulates P53/TP53 activity, P53/TP53 ubiquitination and proteasomal degradation. Acts as a scaffold, binding to both PDPK1 and SRC, thereby allowing SRC to phosphorylate PDPK1 at 'Tyr-9', 'Tyr-373', and 'Tyr-376'. Promotes phosphorylation of NMDA receptors by SRC family members, and thereby contributes to the regulation of NMDA receptor ion channel activity and intracellular Ca (2+) levels. May also regulate potassium ion transport by phosphorylation of potassium channel subunits. Phosphorylates SRC; this increases SRC kinase activity. Phosphorylates ASAP1, NPHP1, KCNA2 and SHC1. Promotes phosphorylation of ASAP2, RHOU and PXN; this requires both SRC and PTK2/PYK2.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	27886584
NCBI GeneID	2185
NCBI Accession #	NP_004094.3
NCBI GenBank Nucleotide #	NM_004103.4
UniProt Accession #	Q14289; Q13475; Q14290; Q16709; Q6PID4; D3DST0
Molecular Weight	115,875 Da
NCBI Official Full Name	protein-tyrosine kinase 2-beta isoform a
NCBI Official Synonym Full Names	protein tyrosine kinase 2 beta
NCBI Official Symbol	PTK2B
NCBI Official Synonym Symbols	PKB; PTK; CAKB; FAK2; PYK2; CADTK; FADK2; RAFTK
NCBI Protein Information	protein-tyrosine kinase 2-beta; FADK 2; CAK-beta; protein kinase B; focal adhesion kinase 2; cell adhesion kinase beta; proline-rich tyrosine kinase 2; calcium-dependent tyrosine kinase; PTK2B protein tyrosine kinase 2 beta; related adhesion focal tyrosin
UniProt Protein Name	Protein-tyrosine kinase 2-beta
UniProt Gene Name	PTK2B
UniProt Synonym Gene Names	FAK2; PYK2; RAFTK; CADTK; CAK-beta; CAKB; FADK 2; RAFTK
UniProt Entry Name	FAK2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/pyk2-antibody-immunohistochemistry-western-blot-elisa/294036>