

Rabbit Smad3 Polyclonal Antibody | anti-Smad3 antibody

Smad3 (phospho Ser208) Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA294064	Rabbit	Human, Mouse, Rat	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Smad3; Polyclonal Antibody; Smad3 (phospho Ser208) Polyclonal Antibody; SMAD3; MADH3; Mothers against decapentaplegic homolog 3; MAD homolog 3; Mad3; Mothers against DPP homolog 3; hMAD-3; JV15-2; SMAD family member 3; SMAD 3; hSMAD3; anti-Smad3 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-Smad3 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-Smad3 antibody

Receptor-regulated SMAD (R-SMAD) that is an intracellular signal transducer and transcriptional modulator activated by TGF-beta (transforming growth factor) and activin type 1 receptor kinases. Binds the TRE element in the promoter region of many genes that are regulated by TGF-beta and, on formation of the SMAD3/SMAD4 complex, activates transcription. Also can form a SMAD3/SMAD4/JUN/FOS complex at the AP-1/SMAD site to regulate TGF-beta-mediated transcription. Has an inhibitory effect on wound healing probably by modulating both growth and migration of primary keratinocytes and by altering the TGF- mediated chemotaxis of monocytes. This effect on wound healing appears to be hormone-sensitive. Regulator of chondrogenesis and osteogenesis and inhibits early healing of bone fractures. Positively regulates PDPK1 kinase activity by stimulating its dissociation from the 14-3-3 protein YWHAQ which acts as a negative regulator.

NCBI AND UNIPROT INFORMATION

NCBI GI #	223029440
NCBI GeneID	4088
NCBI Accession #	NP_001138574.1
NCBI GenBank Nucleotide #	NM_001145102.1
UniProt Accession #	P84022; O09064; O09144; O14510; O35273; Q92940; Q93002; A8K4B6; B7Z4Z5; B7Z6M9; B7Z9Q2; F5H383
Molecular Weight	48,081 Da
NCBI Official Full Name	mothers against decapentaplegic homolog 3 isoform 2
NCBI Official Synonym Full Names	SMAD family member 3
NCBI Official Symbol	SMAD3
NCBI Official Synonym Symbols	LDS3; LDS1C; MADH3; JV15-2; HSPC193; HsT17436
NCBI Protein Information	mothers against decapentaplegic homolog 3; mad3; hMAD-3; hSMAD3; MAD homolog 3; mad homolog JV15-2; mad protein homolog; mothers against DPP homolog 3; SMA- and MAD-related protein 3; SMAD, mothers against DPP homolog 3; MAD, mothers against decapentapleg
UniProt Protein Name	Mothers against decapentaplegic homolog 3
UniProt Gene Name	SMAD3
UniProt Synonym Gene Names	MADH3; MAD homolog 3; Mad3; Mothers against DPP homolog 3; hMAD-3; SMAD 3; Smad3; hSMAD3
UniProt Entry Name	SMAD3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/smad3-antibody-immunohistochemistry-western-blot-elisa/294064>