

Rabbit Hrs Polyclonal Antibody | anti-Hrs antibody

Hrs Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA295026	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS


Hrs; Polyclonal Antibody; Hrs Polyclonal Antibody; HGS; HRS; Hepatocyte growth factor-regulated tyrosine kinase substrate; Protein pp110; anti-Hrs antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-Hrs antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.

WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-Hrs antibody

Involved in intracellular signal transduction mediated by cytokines and growth factors. When associated with STAM, it suppresses DNA signaling upon stimulation by IL-2 and GM-CSF. Could be a direct effector of PI3-kinase in vesicular pathway via early endosomes and may regulate trafficking to early and late endosomes by recruiting clathrin. May concentrate ubiquitinated receptors within clathrin-coated regions. Involved in down-regulation of receptor tyrosine kinase via multivesicular body (MVBs) when complexed with STAM (ESCRT-0 complex). The ESCRT-0 complex binds ubiquitin and acts as sorting machinery that recognizes ubiquitinated receptors and transfers them to further sequential lysosomal sorting/trafficking processes. May contribute to the efficient recruitment of SMADs to the activin receptor complex. Involved in receptor recycling via its association with the CART complex, a multiprotein complex required for efficient transferrin receptor recycling but not for EGFR degradation.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4758528
NCBI GeneID	9146
NCBI Accession #	NP_004703.1
NCBI GenBank Nucleotide #	NM_004712.4
UniProt Accession #	O14964; Q9NR36
Molecular Weight	86,192 Da
NCBI Official Full Name	hepatocyte growth factor-regulated tyrosine kinase substrate
NCBI Official Synonym Full Names	hepatocyte growth factor-regulated tyrosine kinase substrate
NCBI Official Symbol	HGS
NCBI Official Synonym Symbols	HRS
NCBI Protein Information	hepatocyte growth factor-regulated tyrosine kinase substrate; protein pp110; human growth factor-regulated tyrosine kinase substrate
UniProt Protein Name	Hepatocyte growth factor-regulated tyrosine kinase substrate
UniProt Gene Name	HGS
UniProt Synonym Gene Names	HRS
UniProt Entry Name	HGS_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/hrs-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/295026>