

Rabbit HSP10 Polyclonal Antibody | anti-HSP10 antibody

HSP10 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA295035	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

HSP10; Polyclonal Antibody; HSP10 Polyclonal Antibody; HSPE1; 10 kDa heat shock protein; mitochondrial; Hsp10; 10 kDa chaperonin; Chaperonin 10; CPN10; Early-pregnancy factor; EPF; anti-HSP10 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-HSP10 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-HSP10 antibody

Co-chaperonin implicated in mitochondrial protein import and macromolecular assembly. Together with Hsp60, facilitates the correct folding of imported proteins. May also prevent misfolding and promote the refolding and proper assembly of unfolded polypeptides generated under stress conditions in the mitochondrial matrix (PubMed:1346131, PubMed:11422376). The functional units of these chaperonins consist of heptameric rings of the large subunit Hsp60, which function as a back-to-back double ring. In a cyclic reaction, Hsp60 ring complexes bind one unfolded substrate protein per ring, followed by the binding of ATP and association with 2 heptameric rings of the co-chaperonin Hsp10. This leads to sequestration of the substrate protein in the inner cavity of Hsp60 where, for a certain period of time, it can fold undisturbed by other cell components. Synchronous hydrolysis of ATP in all Hsp60 subunits results in the dissociation of the chaperonin rings and the release of ADP and the folded substrate protein (Probable).

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504523
NCBI GeneID	3336
NCBI Accession #	NP_002148.1
NCBI GenBank Nucleotide #	NM_002157.2
UniProt Accession #	P61604; O95421; Q04984; Q53X54; Q9UDH0
Molecular Weight	10,932 Da
NCBI Official Full Name	10 kDa heat shock protein, mitochondrial
NCBI Official Synonym Full Names	heat shock 10kDa protein 1 (chaperonin 10)
NCBI Official Symbol	HSPE1
NCBI Official Synonym Symbols	EPF; CPN10; GROES; HSP10
NCBI Protein Information	10 kDa heat shock protein, mitochondrial; chaperonin 10; 10 kDa chaperonin; early-pregnancy factor; heat shock 10kD protein 1 (chaperonin 10)
UniProt Protein Name	10 kDa heat shock protein, mitochondrial
UniProt Gene Name	HSPE1
UniProt Synonym Gene Names	Hsp10; CPN10; EPF
UniProt Entry Name	CH10_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/hsp10-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/295035>