

Rabbit anti-Human, Mouse MITF Polyclonal Antibody | anti-MITF antibody

MITF Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA295234	Rabbit	Human, Mouse	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MITF; Polyclonal Antibody; MITF Polyclonal Antibody; BHLHE32; Microphthalmia-associated transcription factor; Class E basic helix-loop-helix protein 32; bHLHe32; anti-MITF antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Applicable Applications for anti-MITF antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.
Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-MITF antibody

Transcription factor that regulates the expression of genes with essential roles in cell differentiation, proliferation and survival. Binds to M-boxes (5'-TCATGTG-3') and symmetrical DNA sequences (E-boxes)(5'-CACGTG-3') found in the promoters of target genes, such as BCL2 and tyrosinase (TYR). Plays an important role in melanocyte development by regulating the expression of tyrosinase (TYR) and tyrosinase-related protein 1 (TYRP1). Plays a critical role in the differentiation of various cell types, such as neural crest-derived melanocytes, mast cells, osteoclasts and optic cup-derived retinal pigment epithelium.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557755
NCBI GeneID	4286
NCBI Accession #	NP_000239.1
NCBI GenBank Nucleotide #	NM_000248.3
UniProt Accession #	O75030; Q14841; Q9P2V0; Q9P2V1; Q9P2V2; Q9P2Y8; B4DJL2; D3K197; E9PFN0
Molecular Weight	58,795 Da
NCBI Official Full Name	microphthalmia-associated transcription factor isoform 4
NCBI Official Synonym Full Names	microphthalmia-associated transcription factor
NCBI Official Symbol	MITF
NCBI Official Synonym Symbols	MI; WS2; CMM8; WS2A; bHLHe32
NCBI Protein Information	microphthalmia-associated transcription factor; class E basic helix-loop-helix protein 32
UniProt Protein Name	Microphthalmia-associated transcription factor
UniProt Gene Name	MITF
UniProt Synonym Gene Names	BHLHE32; bHLHe32
UniProt Entry Name	MITF_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/mitf-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/295234>