

Rabbit RANKL Polyclonal Antibody | anti-RANKL antibody

RANKL Polyclonal Antibody

CATALOG # AAA295702	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunofluorescence, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

RANKL; Polyclonal Antibody; RANKL Polyclonal Antibody; TNFSF11; OPGL; TRANCE; Tumor necrosis factor ligand superfamily member 11; Osteoclast differentiation factor; ODF; Osteoprotegerin ligand; Receptor activator of nuclear factor kappa-B ligand; TNF-related activation-induc; anti-RANKL antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-RANKL antibody	IF (Immunofluorescence), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.


WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-RANKL antibody

Cytokine that binds to TNFRSF11B/OPG and to TNFRSF11A/RANK. Osteoclast differentiation and activation factor. Augments the ability of dendritic cells to stimulate naive T-cell proliferation. May be an important regulator of interactions between T-cells and dendritic cells and may play a role in the regulation of the T-cell-dependent immune response. May also play an important role in enhanced bone-resorption in humoral hypercalcemia of malignancy (PubMed:22664871). Induces osteoclastogenesis by activating multiple signaling pathways in osteoclast precursor cells, chief among which is induction of long lasting oscillations in the intracellular concentration of Ca (2+) resulting in the activation of NFATC1, which translocates to the nucleus and induces osteoclast-specific gene transcription to allow differentiation of osteoclasts. During osteoclast differentiation, in a TMEM64 and ATP2A2-dependent manner induces activation of CREB1 and mitochondrial ROS generation necessary for proper osteoclast generation (By similarity).

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507595
NCBI GeneID	8600
NCBI Accession #	NP_003692.1
NCBI GenBank Nucleotide #	NM_003701.3
UniProt Accession #	O14788; O14723; Q96Q17; Q9P2Q3
Molecular Weight	35,478 Da
NCBI Official Full Name	tumor necrosis factor ligand superfamily member 11 isoform 1
NCBI Official Synonym Full Names	tumor necrosis factor (ligand) superfamily, member 11
NCBI Official Symbol	TNFSF11
NCBI Official Synonym Symbols	ODF; OPGL; sOdf; CD254; OPTB2; RANKL; TRANCE; hRANKL2
NCBI Protein Information	tumor necrosis factor ligand superfamily member 11; osteoprotegerin ligand; osteoclast differentiation factor; TNF-related activation-induced cytokine; receptor activator of nuclear factor kappa B ligand; receptor activator of nuclear factor kappa-B ligand
UniProt Protein Name	Tumor necrosis factor ligand superfamily member 11
UniProt Gene Name	TNFSF11
UniProt Synonym Gene Names	OPGL; RANKL; TRANCE; ODF; OPGL; RANKL; TRANCE
UniProt Entry Name	TNF11_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/rankl-antibody-immunofluorescence-western-blot-elisa/295702>