

Rabbit TFIID Polyclonal Antibody | anti-TFIID antibody

TFIID Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA295905	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

TFIID; Polyclonal Antibody; TFIID Polyclonal Antibody; TBP; GTF2D1; TF2D; TATA-box-binding protein; TATA sequence-binding protein; TATA-binding factor; TATA-box factor; Transcription initiation factor TFIID TBP subunit; anti-TFIID antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Applicable Applications for anti-TFIID antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.


IMAGES



IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-TFIID antibody

General transcription factor that functions at the core of the DNA-binding multiprotein factor TFIID (PubMed:2363050, PubMed:2194289, PubMed:9836642, PubMed:27193682). Binding of TFIID to the TATA box is the initial transcriptional step of the pre-initiation complex (PIC), playing a role in the activation of eukaryotic genes transcribed by RNA polymerase II (PubMed:2363050, PubMed:2194289, PubMed:9836642, PubMed:27193682). Component of a BRF2-containing transcription factor complex that regulates transcription mediated by RNA polymerase III (PubMed:26638071). Component of the transcription factor SL1/TIF-IB complex, which is involved in the assembly of the PIC (pre-initiation complex) during RNA polymerase I-dependent transcription (PubMed:15970593). The rate of PIC formation probably is primarily dependent on the rate of association of SL1 with the rDNA promoter. SL1 is involved in stabilization of nucleolar transcription factor 1/UBTF on rDNA.

NCBI AND UNIPROT INFORMATION

NCBI GI #	285026520
NCBI GeneID	6908
NCBI Accession #	NP_001165556.1
NCBI GenBank Nucleotide #	NM_001172085.1
UniProt Accession #	P20226; Q16845; Q6IBM6; Q9UC02; B4E3B3; F5H869
Molecular Weight	37,698 Da
NCBI Official Full Name	TATA-box-binding protein isoform 2
NCBI Official Synonym Full Names	TATA box binding protein
NCBI Official Symbol	TBP
NCBI Official Synonym Symbols	HDL4; GTF2D; SCA17; TFIID; GTF2D1
NCBI Protein Information	TATA-box-binding protein; TATA-box factor; TATA sequence-binding protein; TATA-box binding protein N-terminal domain; transcription initiation factor TFIID TBP subunit
UniProt Protein Name	TATA-box-binding protein
UniProt Gene Name	TBP
UniProt Synonym Gene Names	GTF2D1; TF2D; TFIID
UniProt Entry Name	TBP_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/tfiid-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/295905>