

Rabbit anti-Human MCM3 Polyclonal Antibody | anti-MCM3 antibody

MCM3 Polyclonal Antibody

CATALOG # AAA296169	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

MCM3; Polyclonal Antibody; MCM3 Polyclonal Antibody; DNA replication licensing factor MCM3; DNA polymerase alpha holoenzyme-associated protein P1; P1-MCM3; RLF subunit beta; p102; anti-MCM3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Applicable Applications for anti-MCM3 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.




IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



IHC (Immunohistochemistry)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.



WB (Western Blot)

Dilution: Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.

ADDITIONAL INFORMATION

Related Product Information for anti-MCM3 antibody

Acts as component of the MCM2-7 complex (MCM complex) which is the putative replicative helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. The active ATPase sites in the MCM2-7 ring are formed through the interaction surfaces of two neighboring subunits such that a critical structure of a conserved arginine finger motif is provided in trans relative to the ATP-binding site of the Walker A box of the adjacent subunit. The six ATPase active sites, however, are likely to contribute differentially to the complex helicase activity. Required for DNA replication and cell proliferation.

NCBI AND UNIPROT INFORMATION

NCBI GI #	19857543
NCBI GeneID	4172
UniProt Accession #	P25205; Q92660; Q9BTR3; Q9NUE7; B4DWW4
Molecular Weight	808
NCBI Official Full Name	DNA replication licensing factor MCM3
NCBI Official Synonym Full Names	minichromosome maintenance complex component 3
NCBI Official Symbol	MCM3
NCBI Official Synonym Symbols	HCC5; P1.h; RLFB; P1-MCM3
NCBI Protein Information	DNA replication licensing factor MCM3; p102; RLF subunit beta; hRlf beta subunit; DNA replication factor MCM3; cervical cancer proto-oncogene 5; minichromosome maintenance deficient 3; replication licensing factor, beta subunit; MCM3 minichromosome mainte
UniProt Protein Name	DNA replication licensing factor MCM3
UniProt Gene Name	MCM3
UniProt Entry Name	MCM3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/mcm3-antibody-immunohistochemistry-western-blot-elisa/296169>