

Rabbit anti-Human IRF3 Polyclonal Antibody | anti-IRF3 antibody

IRF3 Polyclonal Antibody

CATALOG # AAA296511	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

IRF3; Polyclonal Antibody; IRF3 Polyclonal Antibody; Interferon regulatory factor 3 (IRF-3); anti-IRF3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Applicable Applications for anti-IRF3 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Notes	ECO:0000303 PubMed:9803267, ECO:0000312 HGNC: HGNC:6118
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 2 weeks. For long term storage store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Dilution: WB 1:500-2000, IHC-p 1:500-200, ELISA 1:10000-20000



IHC (Immunohistochemistry)

Dilution: WB 1:500-2000, IHC-p 1:500-200, ELISA 1:10000-20000



WB (Western Blot)

Dilution: WB 1:500-2000, IHC-p 1:500-200, ELISA 1:10000-20000

ADDITIONAL INFORMATION

Related Product Information for anti-IRF3 antibody	Key transcriptional regulator of type I interferon (IFN)-dependent immune responses which plays a critical role in the innate immune response against DNA and RNA viruses. Regulates the transcription of type I IFN genes (IFN-alpha and IFN-beta) and IFN-stimulated genes (ISG) by binding to an interferon-stimulated response element (ISRE) in their promoters. Acts as a more potent activator of the IFN-beta (IFNB) gene than the IFN-alpha (IFNA) gene and plays a critical role in both the early and late phases of the IFNA/B gene induction. Found in an inactive form in the cytoplasm of uninfected cells and following viral infection, double-stranded RNA (dsRNA), or toll-like receptor (TLR) signaling, is phosphorylated by IKBKE and TBK1 kinases. This induces a conformational change, leading to its dimerization and nuclear localization and association with CREB binding protein (CREBBP) to form dsRNA-activated factor 1 (DRAF1), a complex which activates the transcription of the type I IFN and ISG genes. Can activate distinct gene expression programs in macrophages and can induce significant apoptosis in primary macrophages.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	308199450
NCBI GeneID	3661
NCBI Accession #	NP_001184053.1
NCBI GenBank Nucleotide #	NM_001197124.1
UniProt Accession #	Q14653; Q5FBY1; A8K7L2
Molecular Weight	47,219 Da
NCBI Official Full Name	interferon regulatory factor 3 isoform 4
NCBI Official Synonym Full Names	interferon regulatory factor 3
NCBI Official Symbol	IRF3
NCBI Protein Information	interferon regulatory factor 3
UniProt Protein Name	Interferon regulatory factor 3
UniProt Gene Name	IRF3
UniProt Synonym Gene Names	IRF-3
UniProt Entry Name	IRF3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/irf3-antibody-immunohistochemistry-western-blot-elisa/296511>