

Rabbit anti-Human MAP2K3 Polyclonal Antibody | anti-MAP2K3 antibody

MAP2K3 (Phospho-Thr222) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA301184	Rabbit	Human	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MAP2K3; Polyclonal Antibody; MAP2K3 (Phospho-Thr222) Antibody; MAPKK 3; MEK3; MP2K3; PRKMK3; anti-MAP2K3 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	The antibody detects endogenous levels of MAP2K3 only when phosphorylated at threonine 222.
Purity/Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using
Form/Format	Rabbit IgG in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1.0 mg/ml (varies by lot)
Sequence Length	318
Applicable Applications for anti-MAP2K3 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of threonine 222(A-K-T(p)-M-D) derived from Human MAP2K3.
Modification	Phospho-Thr222
Target Name	MAP2K3
Preparation and Storage	Store at -20 degree C/1 year

IMAGES



IHC (Immunohiostchemistry)
Immunohistochemical analysis of paraffin-embedded human brain tissue using MAP2K3 (Phospho-Thr222) antibody (left) or the same antibody preincubated with blocking peptide (right).



WB (Western Blot)
Western blot analysis of extracts from Jurkat cells treated with serum using MAP2K3 (Phospho-Thr222) Antibody. The lane on the right is treated with the antigen-specific peptide.

ADDITIONAL INFORMATION

Related Product Information for anti-MAP2K3 antibody	The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase is activated by mitogenic and environmental stress, and participates in the MAP kinase-mediated signaling cascade. It phosphorylates and thus activates MAPK14/p38-MAPK. This kinase can be activated by insulin, and is necessary for the expression of glucose transporter. Expression of RAS oncogene is found to result in the accumulation of the active form of this kinase, which thus leads to the constitutive activation of MAPK14, and confers oncogenic transformation of primary cells. Derijard B., Science 267:682-685(1995).Moriguchi T., J. Biol. Chem. 271:26981-26988(1996).The MGC Project Team; Genome Res. 14:2121-2127(2004).
Product Categories/Family for anti-MAP2K3 antibody	Phospho-specific Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	21618347
NCBI GeneID	5606
NCBI Accession #	NP_002747.2
NCBI GenBank Nucleotide #	NM_002756.4
UniProt Accession #	P46734; Q99441; Q9UE71; Q9UE72; B3KSK7
Molecular Weight	39,940 Da
NCBI Official Full Name	dual specificity mitogen-activated protein kinase kinase 3 isoform A
NCBI Official Synonym Full Names	mitogen-activated protein kinase kinase 3
NCBI Official Symbol	MAP2K3
NCBI Official Synonym Symbols	MEK3; MKK3; MAPKK3; PRKMK3; SAPKK2; SAPKK-2
NCBI Protein Information	dual specificity mitogen-activated protein kinase kinase 3
UniProt Protein Name	Dual specificity mitogen-activated protein kinase kinase 3
UniProt Gene Name	MAP2K3
UniProt Synonym Gene Names	MEK3; MKK3; PRKMK3; SKK2; MAP kinase kinase 3; MAPKK 3; MEK 3; SAPK kinase 2; SAPKK-2; SAPKK2
UniProt Entry Name	MP2K3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/map2k3-antibody-immunohistochemistry-western-blot-elisa/301184>