

Rabbit anti-Human, Mouse Topoisomerase I Monoclonal Antibody | anti-TOP1 antibody

Topoisomerase I Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA30133	Rabbit	Human, Mouse	Western Blot, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Flow Cytometry, Functional Assay

PRODUCT SYNONYMS


Topoisomerase I; Monoclonal Antibody; Topoisomerase I Antibody; DNA topoisomerase 1 antibody; DNA topoisomerase I antibody; NUP98 fusion gene antibody; TOP 1 antibody; TOP I antibody; TOP1 antibody; TOP1_HUMAN antibody; TOP1 antibody; Topoisomerase (DNA) I antibody; Topoisomerase 1 antibody; Topoisomerase1 antibody; TopoisomeraseI antibody; Type I DNA topoisomerase antibody; anti-TOP1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Monoclonal
Clone Number	SC69-03
Purity/Purification	ProA affinity purified
Form/Format	1*TBS (pH7.4), 1% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.
Applicable Applications for anti-TOP1 antibody	WB (Western Blot), ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), FCM/FACS (Flow Cytometry)
Application Notes	WB: 1:1000-5000 IHC: 1:50-1:200 ICC: 1:50-1:500 FC/FACS: 1:50-1:100
Antibody Type	Recombinant Antibody
Immunogen	Recombinant protein
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C.




FCM (Flow Cytometry)
Flow cytometric analysis of HepG2 cells with TOP1 antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody




ICC (Immunocytochemistry)
ICC staining TOP1 in 293 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




ICC (Immunocytochemistry)
ICC staining TOP1 in SHG-44 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC (Immunocytochemistry)
ICC staining TOP1 in Hela cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin-embedded mouse brain tissue using anti-TOP1 antibody. Counter stained with hematoxylin.



WB (Western Blot)
Western blot analysis of TOP1 on different lysates using anti-TOP1 antibody at 1/1, 000 dilution. Positive control: Lane 1: HepG2 Lane 2: Jurkat Lane 3: MCF-7

ADDITIONAL INFORMATION

Related Product Information for anti-TOP1 antibody	DNA topoisomerases play essential roles in many DNA metabolic processes including DNA repair. Topoisomerases can introduce DNA damage upon exposure to drugs that stabilize the covalent protein-DNA intermediate of the topoisomerase reaction. Lesions in DNA are also able to trap topoisomerase-DNA intermediates. DNA topoisomerase I (Top1) catalyzes the relaxation of supercoiled DNA by a mechanism of transient DNA strand cleavage characterized by the formation of a phosphotyrosyl bond between the DNA end and active site tyrosine. The antitumor agent camptothecin (CPT) reversibly stabilizes the covalent enzyme-DNA intermediate by inhibiting DNA religation. When a replication fork collides with a DNA Top1 cleavage complex, the covalently bound enzyme must be removed from the DNA 3' end before recombination-dependent replication restart. The tyrosyl-DNA phosphodiesterase Tdp1 and the structure-specific endonuclease Rad1-Rad10 function as primary alternative pathways of Top1 repair in Saccharomyces cerevisiae. In the budding yeast S. cerevisiae, DNA topoisomerases I and II can functionally substitute for each other in removing positive and negative DNA supercoils.
Product Categories/Family for anti-TOP1 antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	11225260
NCBI GeneID	7150
NCBI Accession #	NP_003277.1
NCBI GenBank Nucleotide #	NM_003286.2
UniProt Accession #	P11387; O43256; Q12855; Q12856; Q15610; Q5TFY3; Q9UJN0; A8KA78; E1P5W3
Molecular Weight	90,726 Da
NCBI Official Full Name	DNA topoisomerase 1
NCBI Official Synonym Full Names	topoisomerase (DNA) I
NCBI Official Symbol	TOP1
NCBI Official Synonym Symbols	TOPI
NCBI Protein Information	DNA topoisomerase 1; type I DNA topoisomerase
UniProt Protein Name	DNA topoisomerase 1
UniProt Gene Name	TOP1
UniProt Entry Name	TOP1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/monoclonal/topoisomerase-i-antibody-clone-sc69-03-western-blot-immunocytochemistry-immunofluorescence-immunohistochemistry-flow-cytometry-elisa/30133>