

Rabbit MCM2 Polyclonal Antibody | anti-MCM2 antibody

MCM2 Antibody

CATALOG # AAA301821	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

MCM2; Polyclonal Antibody; MCM2 Antibody; BM28; CDCL1; DNA replication licensing factor MCM2; KIAA0030; MCMD2; anti-MCM2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	The antibody detects endogenous levels of total MCM2 protein.
Purity/Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Form/Format	Rabbit IgG in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1.0 mg/ml (varies by lot)
Sequence Length	904
Applicable Applications for anti-MCM2 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Peptide
Immunogen Description	Synthesized peptide derived from human MCM2.
Target Name	MCM2
Preparation and Storage	Store at -20 degree C


IMAGES



WB (Western Blot)
Western blot analysis of extracts from 3T3 cells (Lane 2), k562 cells (Lane 3) and HepG2 cells (Lane 4), using MCM2 antibody. The lane on the left is treated with synthesized peptide.



WB (Western Blot)
Western blot analysis of extracts from 293 cells, using MCM2 antibody.



IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using MCM2 antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-MCM2 antibody	Acts as component of the MCM2-7 complex (MCM complex) which is the putative replicative helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. The active ATPase sites in the MCM2-7 ring are formed through the interaction surfaces of two neighboring subunits such that a critical structure of a conserved arginine finger motif is provided in trans relative to the ATP-binding site of the Walker A box of the adjacent subunit. The six ATPase active sites, however, are likely to contribute differentially to the complex helicase activity. Required for the entry in S phase and for cell division.H Yan, Genes & Dev., Jun 1991; 5: 944 - 957. Yukio Ishimi, J. Biol. Chem., Nov 2001; 276: 42744 - 42752.Muluye E. Liku, Mol. Biol. Cell, Oct 2005; 16: 5026 - 5039.Daniela S. Dimitrova, J. Cell Biol., Aug 1999; 146: 709.
Product Categories/Family for anti-MCM2 antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	33356547
NCBI GeneID	4171
NCBI Accession #	NP_004517.2
NCBI GenBank Nucleotide #	NM_004526.3
UniProt Accession #	P49736; Q14577; Q15023; Q8N2V1; Q969W7; Q96AE1; Q9BRM7

Molecular Weight	101,896 Da
NCBI Official Full Name	DNA replication licensing factor MCM2
NCBI Official Synonym Full Names	minichromosome maintenance complex component 2
NCBI Official Symbol	MCM2
NCBI Official Synonym Symbols	BM28; CCNL1; CDCL1; cdc19; D3S3194; MITOTIN
NCBI Protein Information	DNA replication licensing factor MCM2
UniProt Protein Name	DNA replication licensing factor MCM2
UniProt Gene Name	MCM2
UniProt Synonym Gene Names	BM28; CCNL1; CDCL1; KIAA0030
UniProt Entry Name	MCM2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/mcm2-antibody-immunohistochemistry-western-blot-elisa/301821>