

Rabbit anti-Human cdc25C Polyclonal Antibody | anti-cdc25C antibody

cdc25C (Phospho-Ser216) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA302025	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot



PRODUCT SYNONYMS

cdc25C; Polyclonal Antibody; cdc25C (Phospho-Ser216) Antibody; CDC25M1; MPIP3; anti-cdc25C antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	The antibody detects endogenous level of cdc25C only when phosphorylated at serine 216.
Purity/Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using
Form/Format	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1.0 mg/ml (varies by lot)
Sequence Length	473
Applicable Applications for anti-cdc25C antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of serine 216 (S-P-S(p)-M-P) derived from Human cdc25C.
Modification	Phospho-Ser216
Target Name	cdc25C
Preparation and Storage	Store at -20 degree C for long term preservation (recommended). Store at 4 degree C for short term use.

IMAGES



IHC (Immunohiostchemistry)
Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using cdc25C(Phospho-Ser216) Antibody (left) or the same antibody preincubated with blocking peptide(right).

IF (Immunofluorescence)
Immunofluorescence staining of methanol-fixed HeLa cells using cdc25C(Phospho-Ser216) Antibody.



WB (Western Blot)
Western blot analysis of extracts from 293 cells untreated or treated with serum using cdc25C(Phospho-Ser216) Antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-cdc25C antibody	cdc25C is highly conserved during evolution and it plays a key role in the regulation of cell division. The encoded protein is a tyrosine phosphatase and belongs to the Cdc25 phosphatase family. It directs dephosphorylation of cyclin B-bound CDC2 and triggers entry into mitosis. It is also thought to suppress p53-induced growth arrest. Multiple alternatively spliced transcript variants of this gene have been described, however, the full-length nature of many of them is not known.
Product Categories/Family for anti-cdc25C antibody	Phospho-specific Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	567316253
NCBI GeneID	995
NCBI Accession #	NP_001274511.1
NCBI GenBank Nucleotide #	NM_001287582.1
UniProt Accession #	P30307; Q96PL3; Q9H168; Q9H2E8; Q9H2E9; Q9H2F1; D3DQB8
Molecular Weight	45,608 Da
NCBI Official Full Name	M-phase inducer phosphatase 3 isoform a
NCBI Official Synonym Full Names	cell division cycle 25C
NCBI Official Symbol	CDC25C

NCBI Official Synonym Symbols	CDC25; PPP1R60
NCBI Protein Information	M-phase inducer phosphatase 3
UniProt Protein Name	M-phase inducer phosphatase 3
UniProt Gene Name	CDC25C
UniProt Entry Name	MPIP3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/cdc25c-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/302025>