

Rabbit anti-Human syk Polyclonal Antibody | anti-SYK antibody

syk (Ab-323) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA303543	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

syk; Polyclonal Antibody; syk (Ab-323) Antibody; Spleen tyrosine kinase; anti-SYK antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	The antibody detects endogenous level of total syk protein.
Purity/Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.
Form/Format	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1.0 mg/ml (varies by lot)
Sequence Length	612
Applicable Applications for anti-SYK antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around aa. 321~325 (N-P-Y-E-P) derived from Human syk.
Target Name	syk
Preparation and Storage	Store at -20 degree C for long term preservation (recommended). Store at 4 degree C for short term use.


IMAGES


IF (Immunofluorescence)

Immunofluorescence staining of methanol-fixed HeLa cells using syk(Ab-323) Antibody.


IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using syk(Ab-323) Antibody (left) or the same antibody preincubated with blocking peptide(right).



WB (Western Blot)

Western blot analysis of extracts from JK cells using syk(Ab-323) Antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-SYK antibody	Positive effector of BCR-stimulated responses. Couples the B-cell antigen receptor (BCR) to the mobilization of calcium ion either through a phosphoinositide 3-kinase-dependent pathway, when not phosphorylated on tyrosines of the linker region, or through a phospholipase C-gamma-dependent pathway, when phosphorylated on Tyr-348 and Tyr-352. Thus the differential phosphorylation of Syk can determine the pathway by which BCR is coupled to the regulation of intracellular calcium ion
Product Categories/Family for anti-SYK antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	205277435
NCBI GeneID	6850
NCBI Accession #	NP_001128524.1
NCBI GenBank Nucleotide #	NM_001135052.3
UniProt Accession #	P43405
Molecular Weight	69,510 Da
NCBI Official Full Name	tyrosine-protein kinase SYK isoform Syk(S)
NCBI Official Synonym Full Names	spleen tyrosine kinase
NCBI Official Symbol	SYK
NCBI Official Synonym Symbols	p72-Syk
NCBI Protein Information	tyrosine-protein kinase SYK
UniProt Protein Name	Tyrosine-protein kinase SYK
UniProt Gene Name	SYK
UniProt Entry Name	KSYK_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/syk-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/303543>