

Rabbit anti-Human CDKN2A Polyclonal Antibody | anti-CDKN2A antibody

CDKN2A Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA304393	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

CDKN2A; Polyclonal Antibody; CDKN2A Antibody; ARF; CDK4I; CDKN2; CMM2; anti-CDKN2A antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	The antibody detects endogenous level of total CDKN2A protein.
Purity/Purification	Antibodies were purified by affinity purification using immunogen.
Form/Format	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1.0 mg/ml (varies by lot)
Sequence Length	156
Applicable Applications for anti-CDKN2A antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Peptide
Immunogen Description	A synthetic peptide of human CDKN2A.
Target Name	CDKN2A
Preparation and Storage	Store at -20 degree C

IMAGES


IF (Immunofluorescence)

Immunofluorescence analysis of A549 cell using CDKN2A antibody. Blue: DAPI for nuclear staining.


WB (Western Blot)

Western blot analysis of extracts of various cell lines, using CDKN2A antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-CDKN2A antibody	The division cycle of eukaryotic cells is regulated by a family of protein kinases known as the cyclin-dependent kinases (CDKs). The sequential activation of individual members of this family and their consequent phosphorylation of critical substrates promotes orderly progression through the cell cycle. It has been reported that CDKN2A binds to CDK4 and inhibits the catalytic activity of the CDK4/cyclin D enzymes. CDKN2A seems to act in a regulatory feedback circuit with CDK4, D-type cyclins and retinoblastoma protein (1). The INK4 (inhibitor of cyclin-dependent kinase 4) family consists of four tumor-suppressor proteins: p15(INK4B), CDKN2A(INK4A), p18(INK4C), and p19(INK4D). While their sequences and structures are highly homologous, they show appreciable differences in conformational flexibility, stability, and aggregation tendency (2). Cell cycle arrest at the G1 checkpoint allows completion of critical macromolecular events prior to S phase. Regulators of the G1 checkpoint include an inhibitor of cyclin-dependent kinase, CDKN2A(INK4); two tumor-suppressor proteins, p53 and RB and cyclin D1. CDKN2A(INK4) is a tumor-suppressor protein and that genetic and epigenetic abnormalities in genes controlling the G1 checkpoint can lead to both escape from senescence and cancer formation (3).
Product Categories/Family for anti-CDKN2A antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	4502749
NCBI GeneID	1029
NCBI Accession #	NP_000068.1
NCBI GenBank Nucleotide #	NM_000077.4
UniProt Accession #	P42771; O95440; Q15191; Q5VVJ5; Q96B52; Q9NP05; A5X2G7; D3DRK1; Q8N726
Molecular Weight	8,731 Da
NCBI Official Full Name	cyclin-dependent kinase inhibitor 2A isoform p16INK4a
NCBI Official Synonym Full Names	cyclin-dependent kinase inhibitor 2A
NCBI Official Symbol	CDKN2A
NCBI Official Synonym Symbols	ARF; MLM; P14; P16; P19; CMM2; INK4; MTS1; TP16; CDK4I; CDKN2; INK4A; MTS-1; P14ARF; P19ARF; P16INK4; P16INK4A; P16-INK4A
NCBI Protein Information	cyclin-dependent kinase inhibitor 2A
UniProt Protein Name	Cyclin-dependent kinase inhibitor 2A
UniProt Gene Name	CDKN2A
UniProt Synonym Gene Names	CDK4I; MTS-1; p16-INK4; p16INK4A
UniProt Entry Name	CDN2A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/cdkn2a-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/304393>