

Rabbit PSAP Polyclonal Antibody | anti-PSAP antibody

PSAP Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA305757	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

PSAP; Polyclonal Antibody; PSAP Antibody; FLJ00245; GLBA; MGC110993; SAP1; anti-PSAP antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	The antibody detects endogenous level of total PSAP protein.
Purity/Purification	Antibodies were purified by affinity purification using immunogen.
Form/Format	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1.0 mg/ml (varies by lot)
Sequence Length	527
Applicable Applications for anti-PSAP antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Recombinant Protein
Immunogen Description	Recombinant protein of human PSAP.
Target Name	PSAP
Preparation and Storage	Store at -20 degree C

IMAGES



IF (Immunofluorescence)

Immunofluorescence analysis of MCF7 cell using PSAP antibody. Blue: DAPI for nuclear staining.



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using PSAP antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-PSAP antibody	The PSAP gene encodes prosaposin, a precursor of four small nonenzymatic glycoproteins termed 'sphingolipid activator proteins' (SAPs) that assist in the lysosomal hydrolysis of sphingolipids. After proteolytic processing of the prosaposin protein, these 4 released polypeptides are functional activators. Saposin A is encoded by residues 60 to 143 of PSAP, saposin B by 195 to 275, saposin C by 311 to 390, and saposin D by 405 to 487. They are four 12-14 kDa heatstable glycoproteins. Saposins A-D localize primarily to the lysosomal compartment where they facilitate the catabolism of glycosphingolipids with short oligosaccharide groups. Saposins A-D are required for the hydrolysis of certain sphingolipids by specific lysosomal hydrolases. (PMID: 2001789) Defects in PSAP are the cause of Gaucher disease, Tay-Sachs disease, and metachromatic leukodystrophy (PubMed: 2060627, PMID: 15773042). This PSAP antibody (10801-1-AP) is expected to recognize both saposin A and B.
Product Categories/Family for anti-PSAP antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	110224476
NCBI GeneID	5660
NCBI Accession #	NP_001035930.1
NCBI GenBank Nucleotide #	NM_001042465.1
UniProt Accession #	P07602; P07292; P15793; P78538; P78541; P78546; P78547; P78558; Q53Y86; Q6IBQ6; Q92739; Q92740
Molecular Weight	58,484 Da
NCBI Official Full Name	prosaposin isoform b preproprotein
NCBI Official Synonym Full Names	prosaposin
NCBI Official Symbol	PSAP
NCBI Official Synonym Symbols	GLBA; SAP1
NCBI Protein Information	prosaposin
UniProt Protein Name	Prosaposin
UniProt Gene Name	PSAP
UniProt Synonym Gene Names	GLBA; SAP1; CSAct; SAP-1; SAP-2
UniProt Entry Name	SAP_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/psap-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/305757>