

Rabbit anti-Human SMC1 Polyclonal Antibody | anti-SMC1 antibody

SMC1 (Ab-957) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA305955	Rabbit	Human	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

SMC1; Polyclonal Antibody; SMC1 (Ab-957) Antibody; SM1A; SMC1A; SMC1L1; SMC1 alpha protein; anti-SMC1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	The antibody detects endogenous level of total SMC1 protein.
Purity/Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.
Form/Format	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1.0 mg/ml (varies by lot)
Sequence Length	1233
Applicable Applications for anti-SMC1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around aa. 955-959 (G-S-S-Q-G) derived from Human SMC1.
Target Name	SMC1
Preparation and Storage	Store at -20 degree C for long term preservation (recommended). Store at 4 degree C for short term use.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using SMC1(Ab-957) Antibody (left) or the same antibody preincubated with blocking peptide(right).



WB (Western Blot)

Western blot analysis of extracts from JK cells using SMC1(Ab-957) Antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-SMC1 antibody	Involved in chromosome cohesion during cell cycle and in DNA repair. Central component of cohesin complex. The cohesin complex is required for the cohesion of sister chromatids after DNA replication. The cohesin complex apparently forms a large proteinaceous ring within which sister chromatids can be trapped. At anaphase, the complex is cleaved and dissociates from chromatin, allowing sister chromatids to segregate. The cohesin complex may also play a role in spindle pole assembly during mitosis. Involved in DNA repair via its interaction with BRCA1 and its related phosphorylation by ATM, or via its phosphorylation by ATR. Works as a downstream effector both in the ATM/NBS1 branch and in the ATR/MSH2 branch of S-phase checkpoint.
Product Categories/Family for anti-SMC1 antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	30581135
NCBI GeneID	8243
NCBI Accession #	NP_006297.2
NCBI GenBank Nucleotide #	NM_006306.3
UniProt Accession #	Q14683; O14995; Q16351; Q2M228
Molecular Weight	143,233 Da
NCBI Official Full Name	structural maintenance of chromosomes protein 1A isoform 1
NCBI Official Synonym Full Names	structural maintenance of chromosomes 1A
NCBI Official Symbol	SMC1A
NCBI Official Synonym Symbols	SMC1; SMCB; CDLS2; SB1.8; SMC1L1; DXS423E; SMC1alpha
NCBI Protein Information	structural maintenance of chromosomes protein 1A
UniProt Protein Name	Structural maintenance of chromosomes protein 1A
UniProt Gene Name	SMC1A
UniProt Synonym Gene Names	DXS423E; KIAA0178; SB1.8; SMC1; SMC1L1; SMC protein 1A; SMC-1-alpha; SMC-1A
UniProt Entry Name	SMC1A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/smc1-antibody-immunohistochemistry-western-blot-elisa/305955>