

Rabbit PRKCA Polyclonal Antibody | anti-PRKCA antibody

PRKCA Antibody

CATALOG # AAA306341	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

PRKCA; Polyclonal Antibody; PRKCA Antibody; PKCA; AAG6; PRKACA; PKC-alpha; anti-PRKCA antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	The antibody detects endogenous level of total PRKCA protein.
Purity/Purification	Antibodies were purified by affinity purification using immunogen.
Form/Format	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1.0 mg/ml (varies by lot)
Sequence Length	672
Applicable Applications for anti-PRKCA antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Peptide
Immunogen Description	A synthetic peptide of human PRKCA.
Target Name	PRKCA
Preparation and Storage	Store at -20 degree C


IMAGES


IHC (Immunohistochemistry)


Immunohistochemical analysis of paraffin-embedded rat lung using PRKCA antibody at dilution of 1:100 (400x lens).


IHC (Immunohistochemistry)


Immunohistochemical analysis of paraffin-embedded mouse lung using PRKCA antibody at dilution of 1:100 (400x lens).



IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin-embedded human rectum using PRKCA antibody at dilution of 1:100 (200x lens).



IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin-embedded Pancreas cancer using PRKCA antibody.



WB (Western Blot)
Western blot analysis of extracts of Jurkat and K562 cells, using PRKCA antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-PRKCA antibody	Activation of protein kinase C (PKC) is one of the earliest events in a cascade that controls a variety of cellular responses, including secretion, gene expression, proliferation, and muscle contraction (1,2). PKC isoforms belong to three groups based on calcium dependency and activators. Classical PKCs are calcium-dependent via their C2 domains and are activated by phosphatidylserine (PS), diacylglycerol (DAG), and phorbol esters (TPA, PMA) through their cysteine-rich C1 domains. Both novel and atypical PKCs are calcium-independent, but only novel PKCs are activated by PS, DAG, and phorbol esters (3-5). Members of these three PKC groups contain a pseudo-substrate or autoinhibitory domain that binds to substrate-binding sites in the catalytic domain to prevent activation in the absence of cofactors or activators. Control of PKC activity is regulated through three distinct phosphorylation events. Phosphorylation at Thr500 in the activation loop, the autophosphorylation site at Thr641, and at carboxy-terminal hydrophobic site Ser660 occurs in vivo (2). Atypical PKC isoforms lack hydrophobic region phosphorylation, which correlates with the presence of glutamic acid rather than the serine or threonine residues found in more typical PKC isoforms. Either the enzyme PDK1 or a close relative is responsible for PKC activation. A recent addition to the PKC superfamily is PKCmu (PKD), which is regulated by DAG and TPA through its C1 domain. PKD is distinguished by the presence of a PH domain and by its unique substrate recognition and Golgi localization (6). PKC-related kinases (PRK) lack the C1 domain and do not respond to DAG or phorbol esters. Phosphatidylinositol lipids activate PRKs and small Rho-family GTPases bind to the homology region 1 (HR1) to regulate PRK kinase activity (7).
Product Categories/Family for anti-PRKCA antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506067
NCBI GeneID	5578
NCBI Accession #	NP_002728.1

NCBI GenBank Nucleotide #	NM_002737.2
UniProt Accession #	P17252 ; Q15137 ; Q32M72 ; Q96RE4 ; B5BU22
Molecular Weight	76,750 Da
NCBI Official Full Name	protein kinase C alpha type
NCBI Official Synonym Full Names	protein kinase C, alpha
NCBI Official Symbol	PRKCA
NCBI Official Synonym Symbols	AAG6; PKCA; PRKACA; PKC-alpha
NCBI Protein Information	protein kinase C alpha type
UniProt Protein Name	Protein kinase C alpha type
UniProt Gene Name	PRKCA
UniProt Synonym Gene Names	PKCA; PRKACA; PKC-A; PKC-alpha
UniProt Entry Name	KPCA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/prkca-antibody-immunohistochemistry-western-blot-elisa/306341>