

Rabbit BRAF Polyclonal Antibody | anti-BRAF antibody

BRAF antibody

CATALOG # AAA307155	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

BRAF; Polyclonal Antibody; BRAF antibody; BRAF1; RAFB1; B-RAF1; FLJ95109; NS7; anti-BRAF antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	The antibody detects endogenous level of total BRAF antibody.
Purity/Purification	Antibodies were purified by affinity purification using immunogen.
Form/Format	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1.0 mg/ml (varies by lot)
Sequence Length	766
Applicable Applications for anti-BRAF antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen Type	Peptide
Immunogen Description	A synthetic peptide of human BRAF.
Target Name	BRAF
Preparation and Storage	Store at -20 degree C

IMAGES


IHC (Immunohistochemistry)

Immunohistochemistry analysis of paraffin-embedded human tonsil tissue using BRAF antibody.


WB (Western Blot)

Western blot analysis of extracts of various cell lines, using BRAF antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-BRAF antibody	BRAF: v-raf murine sarcoma viral oncogene homolog B1, also known as BRAF1; RAFB1; B-RAF1; FLJ95109. Entrez Protein NP_004324. It is the main effectors recruited by GTP-bound Ras to activate the MEK-MAP kinase pathway. B-Raf contains three consensus Akt phosphorylationsites (Ser364, Ser428, and Thr439). B-Raf is a key regulatory molecule of the mitogen-activated protein kinase kinase (MEK), it has a long amino-terminal region,the region is essential for homo-dimerization of B-Raf and hetero-dimerization of B-Raf and c-Raf at the plasma membrane, followed by phosphorylation of Thr118 in the amino-terminal B-Raf-specific region. Notably, in calcium ionophore-stimulated HeLa cells, B-Raf could propagate signals to MEK under the basal level of GTP-Ras. Expression of Raf-B is highly restricted with highestlevels in the cerebrum and testes and defects in braf are involved in a wide range of cancers. The BRAF gene mutation is frequently detected in papillary thyroid carcinoma,melanocytic nevi, primary cutaneous melanomas and colorectal cancers.
Product Categories/Family for anti-BRAF antibody	Total protein Ab

NCBI AND UNIPROT INFORMATION

NCBI GI #	33188459
NCBI GeneID	673
NCBI Accession #	NP_004324.2
NCBI GenBank Nucleotide #	NM_004333.4
UniProt Accession #	P15056; Q13878; Q3MIN6; Q9UDP8; Q9Y6T3; A4D1T4; B6HY61; B6HY62; B6HY63; B6HY64; B6HY65; B6HY66
Molecular Weight	84,437 Da
NCBI Official Full Name	serine/threonine-protein kinase B-raf
NCBI Official Synonym Full Names	B-Raf proto-oncogene, serine/threonine kinase
NCBI Official Symbol	BRAF
NCBI Official Synonym Symbols	NS7; BRAF1; RAFB1; B-RAF1
NCBI Protein Information	serine/threonine-protein kinase B-raf
UniProt Protein Name	Serine/threonine-protein kinase B-raf
UniProt Gene Name	BRAF
UniProt Synonym Gene Names	BRAF1; RAFB1
UniProt Entry Name	BRAF_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.