

Mouse IL-2 ELISA Kit | IL-2 elisa kit

Mouse IL-2 ELISA KIT

CATALOG #
AAA309895

REACTIVITY
Mouse


PRODUCT SYNONYMS

IL-2; Mouse IL-2 ELISA KIT; Mouse IL-2; IL-2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	Natural and recombinant Mouse IL-2 Ligand
Assay Type	Solid Phase Sandwich ELISA
Samples	Cell culture supernatant, serum, plasma (EDTA, citrate, heparin)
Detection Range	0.02-1.0 ng/mL
Sensitivity	7pg/mL
Assay Time	3 hour
Sample Volume	20 uL
Detection Method	Colorimetric
Preparation and Storage	Store at 4 degree C

IMAGES



Standard Curve (Sample)

Representative standard curve for IL-2 ELISA. IL-2 was diluted in serial two-fold steps in Sample Diluent.

ADDITIONAL INFORMATION

Related Product Information for IL-2 elisa kit	Interleukin 2 (IL-2) is a pleiotropic cytokine produced primarily by mitogen- or antigen-activated T lymphocytes (1, 2). Human IL-2 (also known as T-cell growth factor) is produced by T-cells in response to antigenic or mitogenic stimulation. IL-2 is a potent lymphoid cell growth factor which exerts its biological activity primarily on T cells promoting proliferation and maturation.Additionally, IL-2 has been found to stimulate growth and differentiation of B cells, NK cells, LAK cells, monocytes, and oligodendrocytes. IL-2 is involved in treatment of cancers such as melanoma and renal cell cancer. It plays a key role in promoting the clonal expansion of antigen-specific T cells. In addition, IL-2 has also been shown to mediate multiple immune responses on a variety of cell types. The sequence of mouse IL-2 cDNA predicts a 169 amino acid (aa) residue precursor glycoprotein containing a 20 aa residue signal peptide that is cleaved to form the mature protein (3, 4). At the amino acid sequence level, mature mouse IL-2 is approximately 60% identical to human IL-2 (1, 2, 5). Whereas human IL-2 is active on mouse cells, mouse IL-2 is species-specific and is inactive on human cells (6). The gene for IL-2 has been mapped to mouse chromosome 3 (7).The biological effects of IL-2 are mediated by specific cell surface receptor complexes. The functional high-affinity receptor for IL-2 is composed of three distinct polypeptide chains (8, 9). IL-2 stimulates the proliferation of thymocytes; stimulates the proliferation and differentiation of activated B cells; promotes the growth, differentiation and cytotoxic activity of monocytes; induces the growth of natural killer cells and stimulates cytokine production by these cells as well as the cytolytic activity of these cells; enhances the production of lymphocyte-activated killer (LAK) cells; and induces the proliferation and differentiation of oligodendrocytes (1).
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4029234
NCBI GeneID	3558
UniProt Accession #	O95974; P60568
Molecular Weight	4,762 Da
NCBI Official Full Name	IL-2, partial
NCBI Official Synonym Full Names	interleukin 2
NCBI Official Symbol	IL2
NCBI Official Synonym Symbols	IL-2; TCGF; lymphokine
NCBI Protein Information	interleukin-2
UniProt Protein Name	IL-2 protein
UniProt Gene Name	IL-2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/mouse-elisa-kits/il-2/309895>