

Rabbit anti-Human, Mouse NT5C3 Polyclonal Antibody | anti-NT5C3 antibody

NT5C3 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA310902	Rabbit	Human, Mouse	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

NT5C3; Polyclonal Antibody; NT5C3 Antibody; P5N1; UMPH1; Cytosolic 5'-nucleotidase 3A; Cytosolic 5'-nucleotidase 3; Cytosolic 5'-nucleotidase III; cN-III; Pyrimidine 5'-nucleotidase 1; P5'N-1; P5N-1; PN-I; Uridine 5'-monophosphate hydrolase 1; p36; anti-NT5C3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Specificity	Recognizes endogenous levels of NT5C3 protein.
Purity/Purification	Immunogen affinity chromatography
Form/Format	Phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol
Concentration	1 mg/mL (varies by lot)
Sequence Length	297
Applicable Applications for anti-NT5C3 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human NT5C3.
Preparation and Storage	Store at -20 degree C.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of NT5C3 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 C in a hidified chamber. Cells were washed with PBST and incubated with Alexa Fluor 647-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohiostchemistry)

Immunohistochemical analysis of NT5C3 staining in human brain formalin fixed paraffin embedded tissue section. The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of MRPS7 expression in rat tongue whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-NT5C3 antibody	Total protein Ab
---	------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	70608080
NCBI GeneID	51251
NCBI Accession #	NP_001002009.1
NCBI GenBank Nucleotide #	NM_001002009.2
UniProt Accession #	Q9H0P0; Q6IPZ1; Q6NXS6; Q7L3G6; Q9P0P5; Q9UC42; Q9UC43; Q9UC44; Q9UC45; A8K253; B2RAA5; B8ZZC4
Molecular Weight	38 kDa
NCBI Official Full Name	cytosolic 5'-nucleotidase 3A isoform 2
NCBI Official Synonym Full Names	5'-nucleotidase, cytosolic IIIA

NCBI Official Symbol	NT5C3A
NCBI Official Synonym Symbols	p36; PN-I; POMP; PSN1; UMPH; NT5C3; P5N-1; UMPH1; hUMP1; P5'N-1; cN-III
NCBI Protein Information	cytosolic 5'-nucleotidase 3A
UniProt Protein Name	Cytosolic 5'-nucleotidase 3A
UniProt Gene Name	NT5C3A
UniProt Synonym Gene Names	NT5C3; P5N1; UMPH1; 7-methylguanosine nucleotidase; cN-III; P5'N-1; P5N-1; PN-I

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/nt5c3-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/310902>