

Rabbit NFAT3 Polyclonal Antibody | anti-NFAT3 antibody

Phospho-NFAT3 (Ser289) Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA31157	Rabbit	Human, Mouse, RatPredicted: Pig, Bovine, Horse, Sheep, Rabbit, Dog	Western Blot, Immunohistochemistry, ELISA

PRODUCT SYNONYMS

NFAT3; Polyclonal Antibody; Phospho-NFAT3 (Ser289) Antibody; cytoplasmic 4; NF ATc4; NF-AT3; NF-ATc4; NFAC4_HUMAN; NFATc4; Nuclear factor of activated T cells cytoplasmic 4; Nuclear factor of activated T cells cytoplasmic calcineurin dependent 4; Nuclear factor of activated T-cells; T cell transcription factor NFAT3; T-cell transcription factor NFAT3; anti-NFAT3 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat Predicted: Pig, Bovine, Horse, Sheep, Rabbit, Dog
Clonality	Polyclonal
Isotype	IgG
Specificity	Phospho-NFAT3 (Ser289) Antibody detects endogenous levels of NFAT3 only when phosphorylated at Ser289.
Purity/Purification	Purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.
Form/Format	Liquid; Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	964
Applicable Applications for anti-NFAT3 antibody	WB (Western Blot), IHC (Immunohistochemistry), ELISA (Peptide ELISA)
Application Notes	WB: 1:500-1:2000 IHC: 1:50-1:200 ELISA (Peptide): 1:20000-1:40000
Immunogen	A synthesized peptide derived from human NFAT3 around the phosphorylation site of Ser289.
Tissue Specificity	Highly expressed in placenta, lung, kidney, testis and ovary. Weakly expressed in spleen and thymus. Not expressed in peripheral blood lymphocytes. Detected in hippocampus.
Subcellular Location	Nucleus
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.




WB (Western Blot)

Western blot analysis of Phospho-NFAT3 (Ser289) in lysates of HepG2 Ca²⁺ 40uM 30', using Phospho-NFAT3 (Ser289) Antibody.




IHC (Immunohistochemistry)

At 1/100 staining human kidney tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary




IHC (Immunohistochemistry)

At 1/100 staining rat lung tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary




IHC (Immunohistochemistry)

At 1/100 staining rat brain tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary



IHC (Immunohistochemistry)

At 1/100 staining rat heart tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary



IHC (Immunohistochemistry)

At 1/100 staining mouse kidney tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary

ADDITIONAL INFORMATION

Related Product Information for anti-NFAT3 antibody	Description: Plays a role in the inducible expression of cytokine genes in T-cells, especially in the induction of the IL-2 and IL-4. Transcriptionally repressed by estrogen receptors; this inhibition is further enhanced by estrogen. Increases the transcriptional activity of PPARG and has a direct role in adipocyte differentiation. May play an important role in myotube differentiation. May play a critical role in cardiac development and hypertrophy. May play a role in deafferentation-induced apoptosis of sensory neurons. Post Translational Modifications: Phosphorylated by NFATC-kinases; dephosphorylated by calcineurin. Phosphorylated on Ser-168 and Ser-170 by MTOR, IRAK1, MAPK7 and MAPK14, on Ser-213 and Ser-217 by MAPK8 and MAPK9, and on Ser-289 and Ser-344 by RPS6KA3. Phosphorylated by GSK3B. Ubiquitinated, leading to its degradation by the proteasome and reduced transcriptional activity. Ubiquitination and reduction in transcriptional activity can be further facilitated through GSK3B-dependent phosphorylation. Polyubiquitin linkage is mainly through 'Lys-48'. Subunit Structure: Member of the multicomponent NFATC transcription complex that consists of at least two components, a pre-existing cytoplasmic component NFATC2 and an inducible nuclear component NFATC1. Other members such as NFATC4, NFATC3 or members of the activating protein-1 family, MAF, GATA4 and Cbp/p300 can also bind the complex. NFATC proteins bind to DNA as monomers. Interacts with CREBBP, GATA4, IRAK1, MAPK8, MAPK9 and RPS6KA3. Similarity: Rel Similarity Domain (RSD) allows DNA-binding and cooperative interactions with AP1 factors.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	209862845
NCBI GeneID	4776
NCBI Accession #	NP_001129494.1
NCBI GenBank Nucleotide #	NP_001129494.1
UniProt Accession #	Q14934
Molecular Weight	Observed: 95 kDa Predicted: 96 kDa
NCBI Official Full Name	nuclear factor of activated T-cells, cytoplasmic 4 isoform 1
NCBI Official Synonym Full Names	nuclear factor of activated T cells 4
NCBI Official Symbol	NFATC4
NCBI Official Synonym Symbols	NFAT3; NF-AT3; NF-ATC4
NCBI Protein Information	nuclear factor of activated T-cells, cytoplasmic 4
UniProt Protein Name	Nuclear factor of activated T-cells, cytoplasmic 4
UniProt Gene Name	NFATC4
UniProt Synonym Gene Names	NFAT3; NF-ATc4; NFATc4; NF-AT3
UniProt Entry Name	NFAC4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/nfat3-antibody-western-blot-immunohistochemistry-peptide-elisa-elisa/31157>