

Rabbit SLFN11 Polyclonal Antibody | anti-SLFN11 antibody

SLFN11 Antibody


CATALOG # AAA31226	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS ELISA
------------------------------	-----------------------	--	------------------------------

PRODUCT SYNONYMS

SLFN11; Polyclonal Antibody; SLFN11 Antibody; Schlafen family member 11; SLFN8/9; SLN11_HUMAN; anti-SLFN11 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	SLFN11 Antibody detects endogenous levels of total SLFN11.
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin (Thermo Fisher Scientific).
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, pH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol
Concentration	1mg/ml (varies by lot)
Sequence Length	901
Applicable Applications for anti-SLFN11 antibody	ELISA
Application Notes	IF: 1:100-1:500 ICC: 1:100-1:500 WB: 1:500-1:2000 IHC: 1:50-1:200 ELISA(peptide): 1:20,000-1:40,000
Immunogen	A synthesized peptide derived from human SLFN11, corresponding to a region within N-terminal amino acids.
Fragment	Fab fragment
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.




IF (Immunofluorescence)

AAA31226 staining HepG2 cells by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(AAA31226 1:200) and mouse anti-beta tubulin Ab(T0023 1:200) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary antibody. The nuclear counter stain is DAPI(blue).



IHC (Immunohistochemistry-Paraffin)

AAA31226 at 1/100 staining Human colorectal cancer and adjacent normal tissues by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.




IHC (Immunohistochemistry-Paraffin)

AAA31226 at 1/100 staining Human colorectal cancer by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.




IHC (Immunohistochemistry-Paraffin)

AAA31226 at 1/100 staining Mouse lung tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.



IHC (Immunohistochemistry-Paraffin)

AAA31226 at 1/100 staining Mouse testis tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.



IHC (Immunohistochemistry-Paraffin)

AAA31226 at 1/100 staining Rat testis tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-SLFN11 antibody	Function: Inhibitor of DNA replication that promotes cell death in response to DNA damage (PubMed:22927417, PubMed:26658330, PubMed:29395061). Acts as a guardian of the genome by killing cells with defective replication (PubMed:29395061). Persistently blocks stressed replication forks by opening chromatin across replication initiation sites at stressed replication forks, possibly leading to unwind DNA ahead of the MCM helicase and block fork progression, ultimately leading to cell death (PubMed:29395061). Acts independently of ATR (PubMed:29395061). Also acts as an interferon (IFN)-induced antiviral protein which acts as an inhibitor of retrovirus protein synthesis (PubMed:23000900). Specifically abrogates the production of retroviruses such as human immunodeficiency virus 1 (HIV-1) by acting as a specific inhibitor of the synthesis of retroviruses encoded proteins in a codon-usage-dependent manner (PubMed:23000900). Binds to tRNAs and exploits the unique viral codon bias towards A/T nucleotides (PubMed:23000900). The exact inhibition mechanism is unclear: may either sequester tRNAs, prevent their maturation via post-transcriptional processing or may accelerate their deacylation (PubMed:23000900). Does not inhibit reverse transcription, integration or production and nuclear export of viral RNA (PubMed:23000900). Subcellular Location: Nucleus. Chromosome. Note: Recruited to stressed replication forks carrying extended RPA filaments (PubMed:29395061). Recruited to DNA damage sites via interaction with RPA1 (PubMed:26658330, PubMed:29395061). Tissue Specificity: Exhibits a wider expression range in ovarian and colon adenocarcinoma than in their corresponding healthy tissues. Subunit Structure: Interacts with MCM3 (PubMed:29395061). Interacts with DHX9 (PubMed:26625211, PubMed:29395061). Interacts with RPA1 (PubMed:26658330, PubMed:29395061). Similarity: Belongs to the Schlafen family. Subgroup III subfamily.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	157388957
NCBI GeneID	91607
NCBI Accession #	NP_001098057.1
NCBI GenBank Nucleotide #	NP_001098057.1
UniProt Accession #	Q7Z7L1
Molecular Weight	Observed Molecular Weight: (Observed)102kD. Predicted Molecular Weight: (Calculated)103kDa.
NCBI Official Full Name	schlafen family member 11
NCBI Official Synonym Full Names	schlafen family member 11
NCBI Official Symbol	SLFN11
NCBI Official Synonym Symbols	SLFN8/9
NCBI Protein Information	schlafen family member 11
UniProt Protein Name	Schlafen family member 11
UniProt Gene Name	SLFN11
UniProt Entry Name	SLN11_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.