

Rabbit SMAD2 Polyclonal Antibody | anti-SMAD2 antibody

SMAD2 Rabbit Polyclonal Antibody

CATALOG # AAA313264	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

SMAD2; Polyclonal Antibody; SMAD2 Rabbit Polyclonal Antibody; JV18; JV18-1; MADH2; MADR2; hMAD-2; hSMAD2; Smad2; anti-SMAD2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Applicable Applications for anti-SMAD2 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthetic peptide of human Smad2
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.


IMAGES


IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human urothelial carcinoma using Smad2 Rabbit pAb.


WB (Western Blot)

Western blot analysis of extracts of various cell lines, using Smad2 antibody.



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using Smad2 antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-SMAD2 antibody	The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the <i>Drosophila</i> gene 'mothers against decapentaplegic' (Mad) and the <i>C. elegans</i> gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein mediates the signal of the transforming growth factor (TGF)-beta, and thus regulates multiple cellular processes, such as cell proliferation, apoptosis, and differentiation. This protein is recruited to the TGF-beta receptors through its interaction with the SMAD anchor for receptor activation (SARA) protein. In response to TGF-beta signal, this protein is phosphorylated by the TGF-beta receptors. The phosphorylation induces the dissociation of this protein with SARA and the association with the family member SMAD4. The association with SMAD4 is important for the translocation of this protein into the nucleus, where it binds to target promoters and forms a transcription repressor complex with other cofactors. This protein can also be phosphorylated by activin type 1 receptor kinase, and mediates the signal from the activin. Alternatively spliced transcript variants have been observed for this gene.
Product Categories/Family for anti-SMAD2 antibody	Primary Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	51173730
NCBI GeneID	4087
NCBI Accession #	NP_001003652.1
NCBI GenBank Nucleotide #	NM_001003652.3
UniProt Accession #	Q15796
Molecular Weight	52,306 Da
NCBI Official Full Name	mothers against decapentaplegic homolog 2 isoform 1
NCBI Official Synonym Full Names	SMAD family member 2
NCBI Official Symbol	SMAD2
NCBI Official Synonym Symbols	JV18; MADH2; MADR2; JV18-1; hMAD-2; hSMAD2
NCBI Protein Information	mothers against decapentaplegic homolog 2; MAD homolog 2; mother against DPP homolog 2; Sma- and Mad-related protein 2; SMAD, mothers against DPP homolog 2
UniProt Protein Name	Mothers against decapentaplegic homolog 2
UniProt Gene Name	SMAD2
UniProt Synonym Gene Names	MADH2; MADR2; MAD homolog 2; Mothers against DPP homolog 2; hMAD-2; SMAD 2; Smad2; hSMAD2
UniProt Entry Name	SMAD2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/smad2-antibody-immunohistochemistry-western-blot-elisa/313264>