

Rabbit CD40 Polyclonal Antibody | anti-CD40 antibody

CD40 Rabbit Polyclonal Antibody

CATALOG # AAA313387	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

CD40; Polyclonal Antibody; CD40 Rabbit Polyclonal Antibody; Bp50; CDW40; TNFRSF5; p50; anti-CD40 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Applicable Applications for anti-CD40 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A Synthetic peptide of human CD40
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse spleen using CD40 at dilution of 1:200 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat spleen using CD40 at dilution of 1:200 (40x lens).



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded rat lung using CD40 at dilution of 1:200 (40x lens).

ADDITIONAL INFORMATION

Related Product Information for anti-CD40 antibody	This gene is a member of the TNF-receptor superfamily. The encoded protein is a receptor on antigen-presenting cells of the immune system and is essential for mediating a broad variety of immune and inflammatory responses including T cell-dependent immunoglobulin class switching, memory B cell development, and germinal center formation. AT-hook transcription factor AKNA is reported to coordinately regulate the expression of this receptor and its ligand, which may be important for homotypic cell interactions. Adaptor protein TNFR2 interacts with this receptor and serves as a mediator of the signal transduction. The interaction of this receptor and its ligand is found to be necessary for amyloid-beta-induced microglial activation, and thus is thought to be an early event in Alzheimer disease pathogenesis. Mutations affecting this gene are the cause of autosomal recessive hyper-IgM immunodeficiency type 3 (HIGM3). Multiple alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.
Product Categories/Family for anti-CD40 antibody	Primary Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507581
NCBI GeneID	958
NCBI Accession #	NP_001241.1
NCBI GenBank Nucleotide #	NM_001250.4
UniProt Accession #	P25942; Q53GN5; Q5JY15; Q5U007; Q7M4Q8; Q86YK5; Q9BYU0; E1P5S9
Molecular Weight	30,619 Da
NCBI Official Full Name	tumor necrosis factor receptor superfamily member 5 isoform 1
NCBI Official Synonym Full Names	CD40 molecule, TNF receptor superfamily member 5
NCBI Official Symbol	CD40
NCBI Official Synonym Symbols	p50; Bp50; CDW40; TNFRSF5
NCBI Protein Information	tumor necrosis factor receptor superfamily member 5; CD40L receptor; CD40 type II isoform; B cell-associated molecule; B cell surface antigen CD40; B-cell surface antigen CD40; CD40 antigen (TNF receptor superfamily member 5); tumor necrosis factor recept
UniProt Protein Name	Tumor necrosis factor receptor superfamily member 5
UniProt Gene Name	CD40
UniProt Synonym Gene Names	TNFRSF5

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/cd40-antibody-immunohistochemistry-western-blot-elisa/313387>