

Rabbit PIP4K2A Polyclonal Antibody | anti-PIP4K2A antibody

PIP4K2A Rabbit Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA314045	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

PIP4K2A; Polyclonal Antibody; PIP4K2A Rabbit Polyclonal Antibody; PI5P4KA; PIP5K2A; PIP5KII-alpha; PIP5KIIA; PIPK; anti-PIP4K2A antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Applicable Applications for anti-PIP4K2A antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant fusion protein of human PIP4K2A (NP_005019.2).
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of A549 cells using PIP4K2A.




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse lung using PIP4K2A at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat kidney using PIP4K2A at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using PIP4K2A at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Exposure time: 90s.

ADDITIONAL INFORMATION

Related Product Information for anti-PIP4K2A antibody	Phosphatidylinositol-5,4-bisphosphate, the precursor to second messengers of the phosphoinositide signal transduction pathways, is thought to be involved in the regulation of secretion, cell proliferation, differentiation, and motility. The protein encoded by this gene is one of a family of enzymes capable of catalyzing the phosphorylation of phosphatidylinositol-5-phosphate on the fourth hydroxyl of the myo-inositol ring to form phosphatidylinositol-5,4-bisphosphate. The amino acid sequence of this enzyme does not show homology to other kinases, but the recombinant protein does exhibit kinase activity. This gene is a member of the phosphatidylinositol-5-phosphate 4-kinase family.
Product Categories/Family for anti-PIP4K2A antibody	Primary Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	6857820
NCBI GeneID	5305
NCBI Accession #	NP_005019.2
NCBI GenBank Nucleotide #	NM_005028.4
UniProt Accession #	P48426; P53807; Q5VUX3; B0YJ66; B4DGX2; D3DRV1
Molecular Weight	39,802 Da
NCBI Official Full Name	phosphatidylinositol 5-phosphate 4-kinase type-2 alpha
NCBI Official Synonym Full Names	phosphatidylinositol-5-phosphate 4-kinase, type II, alpha
NCBI Official Symbol	PIP4K2A
NCBI Official Synonym Symbols	PIPK; PI5P4KA; PIP5K2A; PIP5KIIA; PIP5KII-alpha
NCBI Protein Information	phosphatidylinositol 5-phosphate 4-kinase type-2 alpha; PIP5KIII; PIP5KIIalpha; PIP4KII-alpha; PI(5)P 4-kinase type II alpha; PtdIns(4)P-5-kinase B isoform; ptdIns(4)P-5-kinase C isoform; diphosphoinositide kinase 2-alpha; ptdIns(5)P-4-kinase isoform 2-al
UniProt Protein Name	Phosphatidylinositol 5-phosphate 4-kinase type-2 alpha
UniProt Gene Name	PIP4K2A
UniProt Synonym Gene Names	PIP5K2; PIP5K2A; PI(5)P 4-kinase type II alpha; PIP4KII-alpha
UniProt Entry Name	PI42A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/pip4k2a-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/314045>